

Human High mobility group protein HMGI-C ELISA Kit | HMGA2 elisa kit

Human High mobility group protein HMGI-C ELISA Kit

CATALOG #
AAA100182

REACTIVITY
Human


PRODUCT SYNONYMS

High mobility group protein HMGI-C; Human High mobility group protein HMGI-C ELISA Kit; HMGA2 (High-mobility group protein HMGI-C)/BABL/HMGIC/LIPO/HMGI-C/STQTL9/high mobility group AT-hook 2/High mobility group AT-hook protein 2/high mobility group protein/high-mobility group (nonhistone chromosomal) protein isoform I-C; HMGA2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Sequence Length	118
Assay Type	Sandwich ELISA, Double Antibody
Samples	Serum, Plasma, Cell Culture Supernatant, cell or tissue lysate, Other liquid samples
Detection Range	0.156-10ng/ml
Sensitivity	0.094ng/ml
Preparation and Storage	Store entire kit at 2-8C for short-term. For longer-term, please store the microplate & standard at -20C, while the remaining reagents can be stored at 2-8C

IMAGES



Standard Curve (Sample)

Typical Data & Standard Curve

Results of a typical standard run of a Human HMGA2 ELISA Kit are shown below. This standard curve was generated at our lab for demonstration purpose only. Each user should obtain their own standard curve as per experiment. (N/A=not applicable)

X	ng/ml	0	0.156	0.313	0.625	1.25	2.5	5	10
Y	OD450	0.036	0.182	0.315	0.570	0.920	1.343	1.701	2.013

Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for HMGA2 elisa kit

Principle of the Assay||This kit was based on sandwich enzyme-linked immune-sorbent assay technology. Anti-Human HMGA2 antibody was pre-coated onto 96-well plates. And the biotin conjugated anti-Human HMGA2 antibody was used as detection antibodies. The standards, test samples and biotin conjugated detection antibody were added to the wells subsequently, and wash with wash buffer. HRP-Streptavidin was added and unbound conjugates were washed away with wash buffer. TMB substrates were used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the Human HMGA2 amount of sample captured in plate. Read the O.D. absorbance at 450nm in a microplate reader, and then the concentration of Human HMGA2can be calculated.

NCBI AND UNIPROT INFORMATION

NCBI GI #	665506020
NCBI GeneID	8091
NCBI Accession #	NP_001287847.1
NCBI GenBank Nucleotide #	NM_001300918.1
Molecular Weight	9,742 Da
NCBI Official Full Name	high mobility group protein HMGI-C isoform c
NCBI Official Synonym Full Names	high mobility group AT-hook 2
NCBI Official Symbol	HMGA2
NCBI Official Synonym Symbols	BABL; LIPO; HMGIC; HMGI-C; STQTL9
NCBI Protein Information	high mobility group protein HMGI-C
UniProt Protein Name	High mobility group protein HMGI-C
UniProt Gene Name	HMGA2
UniProt Synonym Gene Names	HMGIC
UniProt Entry Name	HMGA2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/hmga2-human-elisa-kits/high-mobility-group-protein-hmgi-c/100182>