

Rat RAC-beta serine/threonine-protein kinase ELISA Kit | Akt2 elisa kit

Rat RAC-beta serine/threonine-protein kinase ELISA Kit

CATALOG #
AAA100371

REACTIVITY
Rat


PRODUCT SYNONYMS

RAC-beta serine/threonine-protein kinase; Rat RAC-beta serine/threonine-protein kinase ELISA Kit; Akt2/PKB beta; Akt2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of A-GHR. No significant cross-reactivity or interference between A-GHR and analogues was observed.
Sequence Length	1143
Assay Type	Sandwich ELISA, Double Antibody
Samples	Serum, Plasma, Cell Culture Supernatant, cell or tissue lysate, Other liquid samples
Detection Range	15.625-1000pg/ml
Sensitivity	9.375pg/ml
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level AGHR were tested 20 times on one plate, respectively. Intra-Assay: CV<8%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level AGHR were tested on 3 different plates, 8 replicates in each plate. CV (%) = SD/meanX100. Inter-Assay: CV<10%
Preparation and Storage	Store entire kit at 2-8C for short-term. For longer-term, please store the microplate & standard at -20C, while the remaining reagents can be stored at 2-8C

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for Akt2 elisa kit

Principle of the Assay: This kit was based on sandwich enzyme-linked immune-sorbent assay technology. Anti-A-GHR antibody was pre-coated onto 96-well plates. And the biotin conjugated anti-A-GHR antibody was used as detection antibodies. The standards, test samples and biotin conjugated detection antibody were added to the wells subsequently, and wash with wash buffer. HRP-Streptavidin was added and unbound conjugates were washed away with wash buffer. TMB substrates were used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the A-GHR amount of sample captured in plate. Read the O.D. absorbance at 450nm in a microplate reader, and then the concentration of A-GHR can be calculated.

NCBI AND UNIPROT INFORMATION

NCBI GI #	686642685
NCBI GeneID	20385790
NCBI Accession #	XP_009313788.1
NCBI GenBank Nucleotide #	XM_009315513.1
NCBI Official Full Name	RAC-beta serine/threonine protein kinase
NCBI Official Symbol	DQ04_08311010
NCBI Protein Information	RAC-beta serine/threonine protein kinase

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/akt2-rat-elisa-kits/rac-beta-serine-threonine-protein-kinase/100371>