

Rat Lactoferrin ELISA Kit | LTF/LF elisa kit

Rat Lactoferrin ELISA Kit

CATALOG #
AAA101050

REACTIVITY
Rat

PRODUCT SYNONYMS

Lactoferrin; Rat Lactoferrin ELISA Kit; LTF/LF (Lactoferrin); LTF/LF elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of Lta. No significant cross-reactivity or interference between Lta and analogues was observed.
Sequence Length	475
Assay Type	Sandwich ELISA, Double Antibody
Samples	Serum, Plasma, Cell Culture Supernatant, cell or tissue lysate, Other liquid samples
Detection Range	125-8000pg/ml
Sensitivity	75pg/ml
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Lta were tested 20 times on one plate, respectively. Intra-Assay: CV<8%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Lta were tested on 3 different plates, 8 replicates in each plate. CV (%) = SD/meanX100. Inter-Assay: CV<10%
Preparation and Storage	Store entire kit at 2-8C for short-term. For longer-term, please store the microplate & standard at -20C, while the remaining reagents can be stored at 2-8C

IMAGES

Typical Data & Standard Curve

Results of a typical standard operation of a LTF/LF ELISA Kit are listed below. This standard curve was generated at our lab for demonstration purpose only. Users shall obtain standard curve as per experiment by themselves. (N/A=not applicable)

X	pg/ml	0	31.25	62.5	125	250	500	1000	2000
Y	OD450	0.152	0.374	0.549	0.826	1.197	1.536	1.881	2.193

Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for LTF/LF elisa kit

Principle of the Assay: This kit was based on sandwich enzyme-linked immune-sorbent assay technology. Anti-Lta antibody was pre-coated onto 96-well plates. And the biotin conjugated anti-Lta antibody was used as detection antibodies. The standards, test samples and biotin conjugated detection antibody were added to the wells subsequently, and washed with wash buffer. HRP-Streptavidin was added and unbound conjugates were washed away with wash buffer. TMB substrates were used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the Lta amount of sample captured in plate. Read the O.D. absorbance at 450nm in a microplate reader, and then the concentration of Lta can be calculated.

NCBI AND UNIPROT INFORMATION

NCBI GI #	2104522
NCBI GeneID	4057
Molecular Weight	73,161 Da
NCBI Official Full Name	lactoferrin, partial
NCBI Official Synonym Full Names	lactotransferrin
NCBI Official Symbol	LTF
NCBI Official Synonym Symbols	LF; HLF2; GIG12; HEL110
NCBI Protein Information	lactotransferrin
UniProt Protein Name	Lactotransferrin
UniProt Gene Name	LTF
UniProt Synonym Gene Names	GIG12; LF; Lactoferrin; Lfcin-H
UniProt Entry Name	TRFL_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/ltf-lf-rat-elisa-kits/lactoferrin/101050>