

Mouse coagulation factor II ELISA Kit | F II elisa kit

Mouse coagulation factor II ELISA Kit

CATALOG #
AAA101676

REACTIVITY
Mouse


PRODUCT SYNONYMS

coagulation factor II; Mouse coagulation factor II ELISA Kit; F II/F2/Thrombin/Coagulation Factor II/coagulation factor II (thrombin) receptor-like 2/Coagulation factor II receptor-like 2/Coagulation factor II receptor-like 2 (protease-activated receptor 3)/PAR-3/PAR3protease-activated receptor 3/protease-activated receptor 3/protease-activated receptor-3/PT/serine protease/Thrombin receptor-like 2; F II elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Sequence Length	607
Assay Type	Sandwich ELISA, Double Antibody
Samples	Serum, Plasma, Cell Culture Supernatant, cell or tissue lysate, Other liquid samples
Detection Range	0.156-10ng/ml
Sensitivity	0.094ng/ml
Preparation and Storage	Store entire kit at 2-8C for short-term. For longer-term, please store the microplate & standard at -20C, while the remaining reagents can be stored at 2-8C

IMAGES



Standard Curve (Sample)

Typical Data & Standard Curve

Results of a typical standard run of a Mouse F II ELISA Kit are shown below. This standard curve was generated at our lab for demonstration purpose only. Each user should obtain their own standard curve as per experiment. (N/A-not applicable)

X	ng/ml	0	0.156	0.313	0.625	1.25	2.5	5	10
Y	OD450	0.001	0.072	0.127	0.281	0.462	0.889	1.600	2.144

Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for F II elisa kit

Principle of the Assay||This kit was based on sandwich enzyme-linked immune-sorbent assay technology. anti-Mouse F II antibody was pre-coated onto 96-well plates. And the biotin conjugated anti-Mouse F II antibody was used as detection antibodies. The standards, test samples and biotin conjugated detection antibody were added to the wells subsequently, and wash with wash buffer. HRP-Streptavidin was added and unbound conjugates were washed away with wash buffer. TMB substrates were used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the Mouse F II amount of sample captured in plate. Read the O.D. absorbance at 450nm in a microplate reader, and then the concentration of Mouse F II can be calculated.

NCBI AND UNIPROT INFORMATION

NCBI GI #	147900570
NCBI GeneID	443652
NCBI Accession #	NP_001085292.1
NCBI GenBank Nucleotide #	NM_001091823.1
NCBI Official Full Name	coagulation factor 2 (thrombin)
NCBI Official Synonym Full Names	coagulation factor 2 (thrombin)
NCBI Official Symbol	f2
NCBI Protein Information	coagulation factor 2 (thrombin)

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/f-ii-mouse-elisa-kits/coagulation-factor-ii/101676>