

Rat Mannma Binding Protein/Mannan Binding Lectin ELISA Kit | MBP/MBL elisa kit

Rat Mannma Binding Protein/Mannan Binding Lectin ELISA Kit

CATALOG #
AAA102250

REACTIVITY
Rat


PRODUCT SYNONYMS

Mannma Binding Protein/Mannan Binding Lectin; Rat Mannma Binding Protein/Mannan Binding Lectin ELISA Kit; MBP/MBL/Mbl2/MBL/MBP/MBP1/MBPD/MBL2D/MBP-C/COLEC1/HMBPC/L-MBP/MBL-C/L-MBP/mannose-binding protein C/mannose-binding lectin (protein C) 2; MBP/MBL elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Assay Type	Sandwich ELISA, Double Antibody
Samples	Serum, Plasma, Cell Culture Supernatant, cell or tissue lysate, Other liquid samples
Detection Range	31.25-2000pg/ml
Sensitivity	18.75pg/ml
Preparation and Storage	Store entire kit at 2-8C for short-term. For longer-term, please store the microplate & standard at -20C, while the remaining reagents can be stored at 2-8C

IMAGES



Standard Curve (Sample)

Results of a typical standard run of a Rat MBP ELISA Kit are shown below. This standard curve was generated at our lab for demonstration purpose only. Each user should obtain their own standard curve as per experiment. (N/A=not applicable)

X	pg/ml	0	31.2	62.5	125	250	500	1000	2000
Y	OD450	0.004	0.065	0.139	0.234	0.453	0.837	1.492	2.289

Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for MBP/MBL elisa kit

Principle of the Assay||This kit was based on sandwich enzyme-linked immune-sorbent assay technology. anti-Rat MBP antibody was pre-coated onto 96-well plates. And the biotin conjugated anti-Rat MBP antibody was used as detection antibodies. The standards, test samples and biotin conjugated detection antibody were added to the wells subsequently, and wash with wash buffer. HRP-Streptavidin was added and unbound conjugates were washed away with wash buffer. TMB substrates were used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the Rat MBP amount of sample captured in plate. Read the O.D. absorbance at 450nm in a microplate reader, and then the concentration of Rat MBP can be calculated.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

