

Mouse FKBP52 Monoclonal Antibody | anti-FKBP4 antibody

FKBP52 Antibody: FITC

CATALOG # AAA102898	HOST Mouse	REACTIVITY Canine, Hamster, Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Immunoprecipitation, Western Blot
-------------------------------	----------------------	---	--


PRODUCT SYNONYMS

FKBP52; Monoclonal Antibody; FKBP52 Antibody: FITC; FK506 binding protein 4; FKBP4; FKBP59; HBI; Hsp56; p52; p59; PPlase; Rotamase; T cell FK506 binding protein; anti-FKBP4 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Canine, Hamster, Human, Mouse, Rat
Clonality	Monoclonal
Isotype	IgG1
Clone Number	Hi52C
Form/Format	Protein G Purified
Concentration	1mg/mL (varies by lot)
Sequence Length	459
Applicable Applications for anti-FKBP4 antibody	IHC (Immunohistochemistry), IP (Immunoprecipitation), WB (Western Blot)
Conjugate	FITC
Storage Buffer	PBS, 50% glycerol, 0.09% sodium azide
Preparation and Storage	-20 degree C

IMAGES



WB (Western Blot)

Western Blot analysis of Human Cell lysates showing detection of FKBP52 protein using Mouse Anti-FKBP52 Monoclonal Antibody, Clone Hi52C. Load: 15 ug. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-FKBP52 Monoclonal Antibody at 1.5 ug/mL for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT.



ICC (Immunocytochemistry)

Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-FKBP52 Monoclonal Antibody, Clone Hi52C. Tissue: MCF-7 cells (metastatic mammary gland/breast cell line). Species: Human. Primary Antibody: Mouse Anti-FKBP52 Monoclonal Antibody at 1:1000. Secondary Antibody: APC Goat Anti-Mouse (red). Courtesy of: Tom Ratajczak, Univ. of W. Australia.




WB (Western Blot)

Western Blot analysis of Rat Brain, Heart, Kidney, Liver, Pancreas, Skeletal muscle, Spleen, Testes, Thymus cell lysates showing detection of FKBP52 protein using Mouse Anti-FKBP52 Monoclonal Antibody, Clone Hi52C. Load: 15 ug. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-FKBP52 Monoclonal Antibody at 1.5 ug/mL for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT.



IHC (Immunohistochemistry)

Immunohistochemistry analysis using Mouse Anti-FKBP52 Monoclonal Antibody, Clone Hi52C. Tissue: backskin. Species: Mouse. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-FKBP52 Monoclonal Antibody at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT. Localization: Epidermis.



IHC (Immunohistochemistry)

Immunohistochemistry analysis using Mouse Anti-FKBP52 Monoclonal Antibody, Clone Hi52C. Tissue: prostate tissue (ductal epithelial cells). Species: Human. Primary Antibody: Mouse Anti-FKBP52 Monoclonal Antibody at 1:1000. Courtesy of: David F. Smith, Mayo Clinic, USA.

ADDITIONAL INFORMATION

Related Product Information for anti-FKBP4 antibody	<p>Background Info: Detects an ~52kDa protein representing FKBP52 in cell lysate. Also detects FKBP52 in whole tissue extracts from rat kidney and rat and mouse testes. Heavy chain migrates close to FKBP52 on SDS PAGE.</p> <p>Scientific Background: Hsp90 is crucial to cellular signaling by its regulation of the folding, activity, and stability of a wide range of client proteins. These client protein complexes may also contain one or more cochaperones (1). One class of Hsp90-binding cochaperone is composed of proteins with a characteristic tetratricopeptide repeat (TPR) domain that forms an Hsp90 binding site. Among the TPR cochaperones of Hsp90 are Hop/Sti1, protein phosphatase PP5, and members of both the FK506- and cyclosporin A-binding families of immunophilins (2). FK506-binding protein 51 (FKBP51) and FKBP52 are large molecular weight immunophilins that are part of the mature glucocorticoid receptor (GR) heterocomplex (3). The N terminal domain of each protein binds FK506 and has peptidyl-prolyl isomerase (PPIase) activity that converts prolyl peptide bonds within target proteins from cis- to trans- proline. The C-terminal domains contain the TPR repeats involved in protein-protein interactions with the Hsp90 (4). Although FKBP52 and FKBP51 share ~75% sequence similarity, they affect hormone binding by glucocorticoid receptor in opposing manners and have different Hsp90-binding characteristics (3). FK506 binding protein 51 kDa (FKBP51 or otherwise referred to as FKBP54) has been identified as a progestininducible gene. This protein is predominantly expressed in murine T cells but in humans, it is abundantly expressed in numerous tissues at levels many times higher than FKBP12. The FKBP51 gene is known to be induced by glucocorticoids (5).</p>
Product Categories/Family for anti-FKBP4 antibody	Chaperones ; Heat Shock ; Trafficking
References	1. Cheung-Flynn J., Roberts P.J., Riggs D.L., and Smith D.F. (2003) J. Biol. Chem. 278(19): 17388-17394. 2. Davies T.H., Ning Y.N., and Sanchez E.R. (2002) J Biol. Chem. 277 (7): 4597-4600. 3. Wu, B. et al. (2004) Proc. Natl. Acad. Sci. USA. 101(22): 8348-8353. 4. Denny W.B., Prapapanich V., Smith D.F., and Scammell J.G. (2005) Endocrinology 146(7):3194-3201. 5. Cox M.B. et al. (2007) Molecular Endocrinology. Epub.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503729
NCBI GeneID	2288
NCBI Accession #	NP_002005.1
NCBI GenBank Nucleotide #	NM_002014.3
UniProt Accession #	Q02790 ; Q9UCP1 ; Q9UCV7 ; D3DUQ1
Molecular Weight	51,805 Da
NCBI Official Full Name	peptidyl-prolyl cis-trans isomerase FKBP4

NCBI Official Synonym Full Names	FK506 binding protein 4, 59kDa
NCBI Official Symbol	FKBP4
NCBI Official Synonym Symbols	HBI; p52; Hsp56; FKBP51; FKBP52; FKBP59; PPIase
NCBI Protein Information	peptidyl-prolyl cis-trans isomerase FKBP4; rotamase; HSP binding immunophilin; 51 kDa FK506-binding protein; FK506-binding protein 4 (59kD); T-cell FK506-binding protein, 59kD; peptidylprolyl cis-trans isomerase
UniProt Protein Name	Peptidyl-prolyl cis-trans isomerase FKBP4
UniProt Gene Name	FKBP4
UniProt Synonym Gene Names	FKBP52; PPIase FKBP4; FKBP51; 52 kDa FKBP; FKBP-52; p59; FKBP-4; HBI
UniProt Entry Name	FKBP4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/fkbp52-antibody-clone-hi52c-immunohistochemistry-immunoprecipitation-western-blot-elisa/102898>