Mouse TrpV3 Monoclonal Antibody | anti-Trpv3 antibody

TrpV3 Antibody: PerCP

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA103141	**Mouse**	**Human, Mouse, Rat**	**Immunofluorescence, Immunoprecipitation, Immunocytochemistry, Western Blot**

PRODUCT SYNONYMS

TrpV3; Monoclonal Antibody; TrpV3 Antibody: PerCP; 1110036I10Rik; MGC124324; MGC124325; Transient receptor potential cation channel subfamily V member 3; Trpv3 heat sensitive channel; TRPV3_HUMAN; Vanilloid receptor 3; Vanilloid receptor like 3; Vanilloid receptor-like 3; VRL 3; VRL-3; VRL3; anti-Trpv3 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Isotype	IgG2a
Clone Number	S15-4
Form/Format	Protein G Purified
Concentration	1mg/mL (varies by lot)
Sequence Length	791
Applicable Applications for anti-Trpv3 antibody	IF (Immunofluorescence), IP (Immunoprecipitation), ICC (Immunocytochemistry), WB (Western Blot)
Conjugate	PerCP
Storage Buffer	PBS pH7.4, 50% glycerol, 0.09% sodium azide
Preparation and Storage	-20 degree C


IMAGES



IHC (Immunohistochemisry)
Immunohistochemistry analysis using Mouse Anti-TrpV3 Monoclonal Antibody, Clone S15-4. Tissue: hippocampus. Species: Human. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-TrpV3 Monoclonal Antibody at 1:1000 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT.



WB (Western Blot)
Western Blot analysis of Human Cell lysates showing detection of TrpV3 protein using Mouse Anti-TrpV3 Monoclonal Antibody, Clone S15-4. Load: 15 ug. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-TrpV3 Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT.



IHC (Immunohistochemistry)
Immunohistochemistry analysis using Mouse Anti-TrpV3 Monoclonal Antibody, Clone S15-4. Tissue: backskin. Species: Mouse. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-TrpV3 Monoclonal Antibody at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT.

ADDITIONAL INFORMATION

Related Product Information for anti-Trpv3 antibody	**Background Info:** Detects ~70kDa **Scientific Background:** Ion channels are integral membrane proteins that help establish and control the small voltage gradient across the plasma membrane of living cells by allowing the flow of ions down their electrochemical gradient (1). They are present in the membranes that surround all biological cells because their main function is to regulate the flow of ions across this membrane. Whereas some ion channels permit the passage of ions based on charge, others conduct based on a ionic species, such as sodium or potassium. Furthermore, in some ion channels, the passage is governed by a gate which is controlled by chemical or electrical signals, temperature, or mechanical forces. There are a few main classifications of gated ion channels. There are voltage- gated ion channels, ligand-gated, other gating systems and finally those that are classified differently, having more exotic characteristics. The first are voltage- gated ion channels which open and close in response to membrane potential. These are then separated into sodium, calcium, potassium, proton, transient receptor, and cyclic nucleotide-gated channels; each of which is responsible for a unique role. Ligand-gated ion channels are also known as ionotropic receptors, and they open in response to specific ligand molecules binding to the extracellular domain of the receptor protein. The other gated classifications include activation and inactivation by second messengers, inward-rectifier potassium channels, calcium-activated potassium channels, two-pore-domain potassium channels, light-gated channels, mechano-sensitive ion channels and cyclic nucleotide-gated channels. Finally, the other classifications are based on less normal characteristics such as two-pore channels, and transient receptor potential channels (2). The TRPV3 protein belongs to a family of nonselective cation channels that function in a variety of processes, including temperature sensation and vasoregulation. The thermosensitive members of this family are expressed in subsets of sensory neurons that terminate in the skin, and are activated at distinct physiological temperatures. This channel is activated at temperatures between 22 and 40 degrees C. The gene lies in close proximity to another family member (TRPV1) gene on chromosome 17, and the two encoded proteins are thought to associate with each other to form heteromeric channels (3, 4).
Product Categories/Family for anti-Trpv3 antibody	Ion Channels; Neuroscience
References	1. Hille B. (2001) Ion Channels of Excitable Membranes, 3rd Ed., Sinauer Associated Inc.: Sunderland, MA USA. 2. www.iochannels.org 3. Masamoto Y., Kawabata F., Fushiki T. (2009) Biosci. Biotechnol. Biochem. 73(5): 1021-1027. 4. Xiao R., et al. (2008) J Biol Chem. 283(10): 6162-6174.

NCBI AND UNIPROT INFORMATION

NCBI GI #	429535830
NCBI GeneID	497948
NCBI Accession #	NP_001020928
NCBI GenBank Nucleotide #	NM_001025757.2
UniProt Accession #	Q4QYD9
Molecular Weight	90,273 Da
NCBI Official Full Name	transient receptor potential cation channel subfamily V member 3
NCBI Official Synonym Full Names	transient receptor potential cation channel, subfamily V, member 3
NCBI Official Symbol	Trpv3
NCBI Protein Information	transient receptor potential cation channel subfamily V member 3; heat sensitive channel TRPV3
UniProt Protein Name	Heat sensitive channel TRPV3
UniProt Gene Name	**Trpv3**
UniProt Entry Name	Q4QYD9_RAT

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: https://stage.aaabiotech.com/monoclonal/trpv3-antibody-clone-s15-4-immunofluorescence-immunoprecipitation-immunocytochemistry-western-blot-elisa/103141