

Mouse anti-Mouse, Rat HCN2 Monoclonal Antibody | anti-Hcn2 antibody

HCN2 Antibody: Biotin

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA103187	Mouse	Mouse, Rat	Immunofluorescence, Immunohistochemistry, Immunoprecipitation, Western Blot


PRODUCT SYNONYMS

HCN2; Monoclonal Antibody; HCN2 Antibody: Biotin; BCNG2; HAC1; brain cyclic nucleotide gated channel 2; Potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel; anti-Hcn2 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Mouse, Rat
Clonality	Monoclonal
Isotype	IgG1
Clone Number	S71-37
Form/Format	Protein G Purified
Concentration	1mg/mL (varies by lot)
Sequence Length	863
Applicable Applications for anti-Hcn2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), IP (Immunoprecipitation), WB (Western Blot)
Conjugate	Biotin
Storage Buffer	PBS pH7.4, 50% glycerol, 0.09% sodium azide
Preparation and Storage	-20 degree C

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry analysis using Mouse Anti-HCN2 Monoclonal Antibody, Clone S71-37. Tissue: hippocampus. Species: Human. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-HCN2 Monoclonal Antibody at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT.



WB (Western Blot)

Western Blot analysis of Rat brain membrane lysate showing detection of HCN2 protein using Mouse Anti-HCN2 Monoclonal Antibody, Clone S71-37. Load: 15 ug. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-HCN2 Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT.



IHC (Immunohistochemistry)
Immunohistochemistry analysis using Mouse Anti-HCN2 Monoclonal Antibody, Clone S71-37.
Tissue: frozen brain section. Species: mouse. Fixation: 10% Formalin Solution for 12-24 hours at RT. Primary Antibody: Mouse Anti-HCN2 Monoclonal Antibody at 1:1000 for 1 hour at RT. Secondary Antibody: HRP/DAB Detection System: Biotinylated Goat Anti-Mouse, Streptavidin Peroxidase, DAB Chromogen (brown) for 30 minutes at RT. Counterstain: Mayer Hematoxylin (purple/blue) nuclear stain at 250-500 ul for 5 minutes at RT.

ADDITIONAL INFORMATION

Related Product Information for anti-Hcn2 antibody	<p>Background Info: Detects ~95kDa. No cross-reactivity against HCN1</p> <p>Scientific Background: Hyperpolarization-activated cyclic nucleotide-gated ion channel 2 (HCN2) is an integral membrane protein that helps establish and control the small voltage gradient across the plasma membrane of living cells by allowing the flow of ions down their electrochemical gradient (1). Ion channels are present in the membranes that surround all biological cells because their main function is to regulate the flow of ions across this membrane. Whereas some ion channels permit the passage of ions based on charge, others conduct based on a ionic species, such as sodium or potassium. Furthermore, in some ion channels, the passage is governed by a gate which is controlled by chemical or electrical signals, temperature, or mechanical forces. There are a few main classifications of gated ion channels. There are voltage- gated ion channels, ligand- gated, other gating systems and finally those that are classified differently, having more exotic characteristics. The first are voltage- gated ion channels which open and close in response to membrane potential. These are then separated into sodium, calcium, potassium, proton, transient receptor, and cyclic nucleotide-gated channels; each of which is responsible for a unique role. Ligand-gated ion channels are also known as ionotropic receptors, and they open in response to specific ligand molecules binding to the extracellular domain of the receptor protein. The other gated classifications include activation and inactivation by second messengers, inward-rectifier potassium channels, calcium-activated potassium channels, two-pore-domain potassium channels, light-gated channels, mechano-sensitive ion channels and cyclic nucleotide-gated channels. Finally, the other classifications are based on less normal characteristics such as two-pore channels, and transient receptor potential channels (2). Specifically, hyperpolarization-activated cation channels of the HCN gene family contribute to spontaneous rhythmic activity in both the heart and brain (3).</p>
Product Categories/Family for anti-Hcn2 antibody	Ion Channels ; Neuroscience
References	1. Hille B. (2001) Ion Channels of Excitable Membranes, 3rd Ed., Sinauer Associated Inc.:Sunderland, MA USA. 2. www.iochannels.org 3. Zong X., et al. (2005) J Biol Chem. 280(40): 34224-34232

NCBI AND UNIPROT INFORMATION

NCBI GI #	50878267
NCBI GeneID	114244
NCBI Accession #	NP_446136.1
NCBI GenBank Nucleotide #	NM_053684.1
UniProt Accession #	Q9JKA9; Q6BCT5; Q9QZW6
Molecular Weight	94,920 Da
NCBI Official Full Name	potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 2
NCBI Official Synonym Full Names	hyperpolarization activated cyclic nucleotide-gated potassium channel 2
NCBI Official Symbol	Hcn2
NCBI Protein Information	potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 2
UniProt Protein Name	Potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 2

UniProt Gene Name	Hcn2
UniProt Entry Name	HCN2_RAT

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/hcn2-antibody-clone-s71-37-immunofluorescence-immunohistochemistry-immunoprecipitation-western-blot-elisa/103187>