Mouse HCN4 Monoclonal Antibody | anti-Hcn4 antibody

HCN4 Antibody: PerCP

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA103220	**Mouse**	**Human, Mouse, Rat**	**Immunofluorescence, Immunohistochemistry, Immunocytochemistry, Western Blot**

PRODUCT SYNONYMS

HCN4; Monoclonal Antibody; HCN4 Antibody: PerCP; Hyperpolarization activated cyclic nucleotide gated potassium channel 4; HCN 4; Potassium/sodium hyperpolarization activated cyclic nucleotide gated channel 4; anti-Hcn4 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Isotype	IgG1
Clone Number	S114-10
Form/Format	Protein G Purified
Concentration	1mg/mL (varies by lot)
Sequence Length	1198
Applicable Applications for anti-Hcn4 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ICC (Immunocytochemistry), WB (Western Blot)
Conjugate	PerCP
Storage Buffer	PBS pH7.4, 50% glycerol, 0.09% sodium azide
Preparation and Storage	-20 degree C


IMAGES



IHC (Immunohistochemisry)
Immunohistochemistry analysis using Mouse Anti-HCN4 Monoclonal Antibody, Clone S114-10. Tissue: hippocampus. Species: Human. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-HCN4 Monoclonal Antibody at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT.



WB (Western Blot)
Western Blot analysis of Human T-HEK cell lysate showing detection of HCN4 protein using Mouse Anti-HCN4 Monoclonal Antibody, Clone S114-10. Load: 15 ug. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-HCN4 Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT.



IHC (Immunohistochemistry)
Immunohistochemistry analysis using Mouse Anti-HCN4 Monoclonal Antibody, Clone S114-10. Tissue: frozen brain section. Species: mouse. Fixation: 10% Formalin Solution for 12-24 hours at RT. Primary Antibody: Mouse Anti-HCN4 Monoclonal Antibody at 1:1000 for 1 hour at RT. Secondary Antibody: HRP/DAB Detection System: Biotinylated Goat Anti-Mouse, Streptavidin Peroxidase, DAB Chromogen (brown) for 30 minutes at RT. Counterstain: Mayer Hematoxylin (purple/blue) nuclear stain at 250-500 ul for 5 minutes at RT.

ADDITIONAL INFORMATION

Related Product Information for anti-Hcn4 antibody	**Background Info:** Detects ~130kDa. No cross-reactivity against other HCNs **Scientific Background:** Ion channels are integral membrane proteins that help establish and control the small voltage gradient across the plasma membrane of living cells by allowing the flow of ions down their electrochemical gradient (1). They are present in the membranes that surround all biological cells because their main function is to regulate the flow of ions across this membrane. Whereas some ion channels permit the passage of ions based on charge, others conduct based on a ionic species, such as sodium or potassium. Furthermore, in some ion channels, the passage is governed by a gate which is controlled by chemical or electrical signals, temperature, or mechanical forces. There are a few main classifications of gated ion channels. There are voltage- gated ion channels, ligand-gated, other gating systems and finally those that are classified differently, having more exotic characteristics. The first are voltage- gated ion channels which open and close in response to membrane potential. These are then separated into sodium, calcium, potassium, proton, transient receptor, and cyclic nucleotide-gated channels; each of which is responsible for a unique role. Ligand-gated ion channels are also known as ionotropic receptors, and they open in response to specific ligand molecules binding to the extracellular domain of the receptor protein. The other gated classifications include activation and inactivation by second messengers, inward-rectifier potassium channels, calcium-activated potassium channels, two-pore-domain potassium channels, light-gated channels, mechano-sensitive ion channels and cyclic nucleotide-gated channels. Finally, the other classifications are based on less normal characteristics such as two-pore channels, and transient receptor potential channels (2). Specifically, hyperpolarization-activated cation channels of the HCN gene family contribute to spontaneous rhythmic activity in both the heart and brain (3).
Product Categories/Family for anti-Hcn4 antibody	Ion Channels; Neuroscience
References	1. Hille B. (2001) Ion Channels of Excitable Membranes, 3rd Ed., Sinauer Associated Inc.:Sunderland, MA USA. 2. www.iochannels.org 3. Zong X., et al. (2005) J Biol Chem. 280(40): 34224-34232

NCBI AND UNIPROT INFORMATION

NCBI GI #	25742571
NCBI GeneID	59266
NCBI Accession #	NP_067690.1
NCBI GenBank Nucleotide #	NM_021658.1
UniProt Accession #	Q9JKA7; Q9QZW4
Molecular Weight	128,761 Da
NCBI Official Full Name	potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 4
NCBI Official Synonym Full Names	hyperpolarization activated cyclic nucleotide-gated potassium channel 4
NCBI Official Symbol	Hcn4
NCBI Protein Information	potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 4; hyperpolarization-activated, cyclic nucleotide-gated K+ 4
UniProt Protein Name	Potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 4

UniProt Gene Name	**Hcn4**
UniProt Entry Name	HCN4_RAT

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: https://stage.aaabiotech.com/monoclonal/hcn4-antibody-clone-s114-10-immunofluorescence-immunohistochemistry-immunocytochemistry-western-blot-elisa/103220