

Mouse Cavbeta2 Monoclonal Antibody | anti-Cacnb2 antibody

CaVbeta2 Antibody: PerCP

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA103389	Mouse	Human, Mouse, Rat	Immunohistochemistry, Immunoprecipitation, Western Blot


PRODUCT SYNONYMS

Cavbeta2; Monoclonal Antibody; CaVbeta2 Antibody: PerCP; Cacnb2; cacnb2; Voltage-dependent L-type calcium channel subunit beta-2; CAB2; anti-Cacnb2 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Isotype	IgG1
Clone Number	S8b-1
Form/Format	Protein G Purified
Concentration	1mg/mL (varies by lot)
Sequence Length	604
Applicable Applications for anti-Cacnb2 antibody	IHC (Immunohistochemistry), IP (Immunoprecipitation), WB (Western Blot)
Conjugate	PerCP
Storage Buffer	PBS pH7.4, 50% glycerol, 0.09% sodium azide
Preparation and Storage	-20 degree C

IMAGES




ICC (Immunocytochemistry)

Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Cav Beta2 Calcium Channel Monoclonal Antibody, Clone S8b-1. Tissue: HaCaT cells. Species: Human. Fixation: Cold 100% methanol for 10 minutes at -20 degree C. Primary Antibody: Mouse Anti-Cav Beta2 Calcium Channel Monoclonal Antibody at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT. Localization: All cells positive. Bright dottiness located throughout cytoplasm and in nuclei.



WB (Western Blot)

Western Blot analysis of Human Cell line lysates showing detection of Cav Beta2 Calcium Channel protein using Mouse Anti-Cav Beta2 Calcium Channel Monoclonal Antibody, Clone S8b-1. Load: 15 ug. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-Cav Beta2 Calcium Channel Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT.



IHC (Immunohistochemistry)

Immunohistochemistry analysis using Mouse Anti-Cav Beta2 Calcium Channel Monoclonal Antibody, Clone S8b-1. Tissue: backskin. Species: Mouse. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-Cav Beta2 Calcium Channel Monoclonal Antibody at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT. Localization: All nuclei. Some nuclei also staining higher up in epidermis.

ADDITIONAL INFORMATION

Related Product Information for anti-Cacnb2 antibody	Background Info: Detects ~78 kDa. No cross reactivity against Cavbeta1, Cavbeta3, Cavbeta4 Scientific Background: Cav Beat subunits are involved in the transport of the pore forming alpha1 subunit to the plasma membrane (3). They also shield an ER Retention signal on the alpha1 subunit, thereby guiding the pore-forming subunit to the target membrane (4, 5). They also determine the biophysical properties of the calcium channel (5).
Product Categories/Family for anti-Cacnb2 antibody	Ion Channels ; Neuroscience
References	1. Hille B. (2001) Ion Channels of Excitable Membranes, 3rd Ed., Sinauer Associated Inc.: Sunderland, MA USA. 2. www.iochannels.org 3. Dolphin A.C. (2003) J Bioenerg. Biomembr. 35: 599-620. 4. Bichet D., et al. (2000) Neuron. 25: 177-190. 5. Xie M., et al. (2007) J Cell Biol. 178(3): 489-502.

NCBI AND UNIPROT INFORMATION

NCBI GI #	16758716
NCBI GeneID	116600
NCBI Accession #	NP_446303
NCBI GenBank Nucleotide #	NM_053851.1
UniProt Accession #	Q8VGC3; Q811Q6; Q811Q7; Q91ZJ8; Q9QUU7
Molecular Weight	68,066 Da
NCBI Official Full Name	voltage-dependent L-type calcium channel subunit beta-2
NCBI Official Synonym Full Names	calcium channel, voltage-dependent, beta 2 subunit
NCBI Official Symbol	Cacnb2
NCBI Official Synonym Symbols	Cacnlb2
NCBI Protein Information	voltage-dependent L-type calcium channel subunit beta-2; CAB2; calcium channel voltage-dependent subunit beta 2
UniProt Protein Name	Voltage-dependent L-type calcium channel subunit beta-2
UniProt Gene Name	Cacnb2
UniProt Synonym Gene Names	Cacnlb2; CAB2
UniProt Entry Name	CACB2_RAT

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.