

Mouse 4-Hydroxynonenal Monoclonal Antibody | anti-4-HNE antibody

4-Hydroxynonenal Antibody, Clone 12F7: ATTO 488


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA104042	Mouse	Species Independent	ELISA, Immunofluorescence, Immunocytochemistry, Western Blot

PRODUCT SYNONYMS

4-Hydroxynonenal; Monoclonal Antibody; 4-Hydroxynonenal Antibody; Clone 12F7: ATTO 488; Mouse Anti-4-Hydroxynonenal (4-HNE) Monoclonal IgG1; 4-Hydroxynonenal Antibody: ATTO 488; 4-Hydroxy Nonenal (4-HNE) Antibody; 4-Hydroxy Nonenal Antibody; 4-HNE Antibody; 4-hydroxy Hexenal Antibody; HNE Antibody; 4-hydroxy-2-nonenal Antibody; anti-4-HNE antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Species Independent
Clonality	Monoclonal
Isotype	IgG1
Clone Number	12F7
Specificity	Specific for 4-Hydroxynonenal (4-HNE) modified proteins. Does not detect free 4-Hydroxynonenal. Does not cross-react with 4-Hydroxy-2-hexenal, Acrolein, Crotonaldehyde, Hexanoyl Lysine, Malondialdehyde, or Methylglyoxal modified proteins.
Purity/Purification	Protein G Purified
Concentration	1 mg/ml (varies by lot)
Applicable Applications for anti-4-HNE antibody	ELISA, IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Target	4-Hydroxynonenal
Immunogen	Synthetic 4-Hydroxynonenal modified Keyhole Limpet Hemocyanin (KLH).
Conjugate	ATTO 488
Storage Buffer	PBS pH 7.4, 50% glycerol, 0.09% Sodium Azide
Preparation and Storage	Store at -20 degree C.



WB (Western Blot)

Western Blot analysis of 4-hydroxy-nonenal-BSA Conjugate showing detection of 67 kDa 4-hydroxy-nonenal protein using Mouse Anti-4-hydroxy-nonenal Monoclonal Antibody, Clone 12F7. Lane 1: Molecular Weight Ladder (MW). Lane 2: BSA (0.5 ug). Lane 3: 4-hydroxyl nonenal-BSA (0.5 ug). Lane 4: 4-hydroxy nonenal-BSA (2.0 ug). Lane 5: 4-hydroxy-2-hexenal (0.5 ug). Lane 6: 4-hydroxy-2-hexenal (2.0 ug). Block: 5% Skim Milk in TBST. Primary Antibody: Mouse Anti-4-hydroxy-nonenal Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:2000 for 60 min at RT. Color Development: ECL solution for 5 min in RT. Predicted/Observed Size: 67 kDa.



ICC (Immunocytochemistry)

Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-4-Hydroxynonenal Monoclonal Antibody, Clone 12F7. Tissue: Embryonic kidney epithelial cell line (HEK293). Species: Human. Fixation: 5% Formaldehyde for 5 min. Primary Antibody: Mouse Anti-4-Hydroxynonenal Monoclonal Antibody at 1:50 for 30-60 min at RT. Secondary Antibody: Goat Anti-Mouse Alexa Fluor 488 at 1:1500 for 30-60 min at RT. Counterstain: Phalloidin Alexa Fluor 633 F-Actin stain; DAPI (blue) nuclear stain at 1:250, 1:50000 for 30-60 min at RT. Magnification: 20X (2X Zoom). (A,C,E,G) - Untreated. (B,D,F,H) - Cells cultured overnight with 50 uM H2O2. (A,B) DAPI (blue) nuclear stain. (C,D) Phalloidin Alex Fluor 633 F-Actin stain. (E,F) 4-Hydroxynonenal Antibody. (G,H) Composite. Courtesy of: Dr. Robert Burke, University of Victoria.

ADDITIONAL INFORMATION

Product Categories/Family for anti-4-HNE antibody	Antibodies; Cancer; Oxidative Stress; Lipid peroxidation
---	--

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/4-hydroxynonenal-antibody-clone-12f7-immunofluorescence-immunocytochemistry-western-blot-elisa/104042>