

## Mouse Malondialdehyde Monoclonal Antibody | anti-MDA antibody

Malondialdehyde Antibody, Clone 6H6: ATTO 390

| CATALOG # | HOST  | REACTIVITY          | APPLICATIONS                                                                                   |
|-----------|-------|---------------------|------------------------------------------------------------------------------------------------|
| AAA104068 | Mouse | Species Independent | ELISA, Flow Cytometry, Functional Assay, Immunofluorescence, Immunocytochemistry, Western Blot |


### PRODUCT SYNONYMS

Malondialdehyde; Monoclonal Antibody; Malondialdehyde Antibody; Clone 6H6: ATTO 390; Mouse Anti-Malondialdehyde (MDA) Monoclonal IgG1; Malondialdehyde Antibody: ATTO 390; MDA Antibody; Malondialdehyde (MDA) Antibody; Malonic aldehyde Antibody Propanedial Antibody; 1; 3-Propanedial Antibody; Malonaldehyde Antibody; anti-MDA antibody

### PRODUCT SPECIFICATIONS


|                                               |                                                                                                                                                                                                                                          |
|-----------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Host                                          | Mouse                                                                                                                                                                                                                                    |
| Reactivity                                    | Species Independent                                                                                                                                                                                                                      |
| Clonality                                     | Monoclonal                                                                                                                                                                                                                               |
| Isotype                                       | IgG1                                                                                                                                                                                                                                     |
| Clone Number                                  | 6H6                                                                                                                                                                                                                                      |
| Specificity                                   | Specific for Malondialdehyde conjugated proteins. Does not detect free Malondialdehyde. Does not cross-react with Acrolein, Crotonaldehyde, Hexanoyl Lysine, 4-Hydroxy-2-hexenal, 4-Hydroxy nonenal, or Methylglyoxal modified proteins. |
| Purity/Purification                           | Protein G Purified                                                                                                                                                                                                                       |
| Concentration                                 | 1 mg/ml (varies by lot)                                                                                                                                                                                                                  |
| Applicable Applications for anti-MDA antibody | ELISA, FCM/FACS (Flow Cytometry), IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)                                                                                                                                  |
| Target                                        | Malondialdehyde                                                                                                                                                                                                                          |
| Immunogen                                     | Synthetic Malondialdehyde modified Keyhole Limpet Hemocyanin (KLH).                                                                                                                                                                      |
| Conjugate                                     | ATTO 390                                                                                                                                                                                                                                 |
| Storage Buffer                                | PBS pH 7.4, 50% glycerol, 0.09% Sodium Azide                                                                                                                                                                                             |
| Preparation and Storage                       | Store at -20 degree C.                                                                                                                                                                                                                   |

### IMAGES



#### FCM/FACS (Flow Cytometry)

Flow Cytometry analysis using Mouse Anti-Malondialdehyde Monoclonal Antibody, Clone 6H6. Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 90% Methanol. Primary Antibody: Mouse Anti-Malondialdehyde Monoclonal Antibody at 1:50 for 30 min on ice. Secondary Antibody: Goat Anti-Mouse: PE at 1:100 for 20 min at RT. Isotype Control: Non Specific IgG. Cells were subject to oxidative stress by treating with 250 uM H2O2 for 24 hours.



#### WB (Western Blot)

Western Blot analysis of Malondialdehyde-BSA Conjugate showing detection of 67 kDa Malondialdehyde protein using Mouse Anti-Malondialdehyde Monoclonal Antibody, Clone 6H6. Lane 1: Molecular Weight Ladder (MW). Lane 2: Malondialdehyde-BSA (0.5 ug). Lane 3: Malondialdehyde-BSA (2.0 ug). Lane 4: BSA (0.5 ug). Lane 5: BSA (2.0 ug) . Block: 5% Skim Milk in TBST. Primary Antibody: Mouse Anti-Malondialdehyde Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:2000 for 60 min at RT. Color Development: ECL solution for 5 min in RT. Predicted/Observed Size: 67 kDa.



#### ICC (Immunocytochemistry)

Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Malondialdehyde Monoclonal Antibody, Clone 6H6. Tissue: Embryonic kidney epithelial cell line (HEK293). Species: Human. Fixation: 5% Formaldehyde for 5 min. Primary Antibody: Mouse Anti-Malondialdehyde Monoclonal Antibody at 1:50 for 30-60 min at RT. Secondary Antibody: Goat Anti-Mouse Alexa Fluor 488 at 1:1500 for 30-60 min at RT. Counterstain: Phalloidin Alexa Fluor 633 F-Actin stain; DAPI (blue) nuclear stain at 1:250, 1:50000 for 30-60 min at RT. Magnification: 20X (2X Zoom). (A,C,E,G) - Untreated. (B,D,F,H) - Cells cultured overnight with 50 uM H2O2. (A,B) DAPI (blue) nuclear stain. (C,D) Phalloidin Alex Fluor 633 F-Actin stain. (E,F) Malondialdehyde Antibody. (G,H) Composite. Courtesy of: Dr. Robert Burke, University of Victoria.

### ADDITIONAL INFORMATION

|                                                 |                                                                                                                |
|-------------------------------------------------|----------------------------------------------------------------------------------------------------------------|
| Product Categories/Family for anti-MDA antibody | Antibodies; Cancer; Oxidative Stress; Lipid peroxidation; Neuroscience; Neurodegeneration; Alzheimer's Disease |
|-------------------------------------------------|----------------------------------------------------------------------------------------------------------------|

**For Research Use Only.** Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | [contact@aaabiotech.com](mailto:contact@aaabiotech.com)  
Online datasheet: <https://stage.aaabiotech.com/monoclonal/malondialdehyde-antibody-clone-6h6-flow-cytometry-immunofluorescence-immunocytochemistry-western-blot-elisa/104068>