

Mouse Malondialdehyde Monoclonal Antibody | anti-MDA antibody

Malondialdehyde Antibody, Clone 11E3: RPE

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA104084	Mouse	Species Independent	ELISA, Immunofluorescence, Immunocytochemistry, Western Blot


PRODUCT SYNONYMS

Malondialdehyde; Monoclonal Antibody; Malondialdehyde Antibody; Clone 11E3: RPE; Mouse Anti-Malondialdehyde (MDA) Monoclonal IgG1; Malondialdehyde Antibody: RPE; MDA Antibody; Malondialdehyde (MDA) Antibody; Malonic aldehyde Antibody Propanedial Antibody; 1; 3-Propanedial Antibody; Malonaldehyde Antibody; anti-MDA antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Species Independent
Clonality	Monoclonal
Isotype	IgG1
Clone Number	11E3
Specificity	Specific for Malondialdehyde conjugated proteins. Does not detect free Malondialdehyde. Does not cross-react with Acrolein, Crotonaldehyde, Hexanoyl Lysine, 4-Hydroxy-2-hexenal, 4-Hydroxy nonenal, or Methylglyoxal modified proteins.
Purity/Purification	Protein G Purified
Concentration	1 mg/ml (varies by lot)
Applicable Applications for anti-MDA antibody	ELISA, IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Target	Malondialdehyde
Immunogen	Synthetic Malondialdehyde modified Keyhole Limpet Hemocyanin (KLH).
Conjugate	RPE
Storage Buffer	PBS pH 7.4, 50% glycerol, 0.09% Sodium Azide
Preparation and Storage	Store at -20 degree C.

IMAGES



WB (Western Blot)

Western Blot analysis of Human Cervical cancer cell line (HeLa) lysate showing detection of Malondialdehyde protein using Mouse Anti-Malondialdehyde Monoclonal Antibody, Clone 11E3. Lane 1: Molecular Weight Ladder (MW). Lane 2: HeLa cell lysate. Lane 3: H₂O₂ treated HeLa cell lysate. Load: 12 ug. Block: 5% Skim Milk in TBST. Primary Antibody: Mouse Anti-Malondialdehyde Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:2000 for 60 min at RT. Color Development: ECL solution for 5 min in RT.



WB (Western Blot)

Western Blot analysis of Malondialdehyde-BSA Conjugate showing detection of 67 kDa Malondialdehyde protein using Mouse Anti-Malondialdehyde Monoclonal Antibody, Clone 11E3. Lane 1: Molecular Weight Ladder (MW). Lane 2: Malondialdehyde-BSA (0.5 ug). Lane 3: Malondialdehyde-BSA (2.0 ug). Lane 4: BSA (0.5 ug). Lane 5: BSA (2.0 ug). Block: 5% Skim Milk in TBST. Primary Antibody: Mouse Anti-Malondialdehyde Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:2000 for 60 min at RT. Color Development: ECL solution for 5 min in RT. Predicted/Observed Size: 67 kDa.



ICC (Immunocytochemistry)

Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Malondialdehyde Monoclonal Antibody, Clone 11E3. Tissue: Embryonic kidney epithelial cell line (HEK293). Species: Human. Fixation: 5% Formaldehyde for 5 min. Primary Antibody: Mouse Anti-Malondialdehyde Monoclonal Antibody at 1:50 for 30-60 min at RT. Secondary Antibody: Goat Anti-Mouse Alexa Fluor 488 at 1:1500 for 30-60 min at RT. Counterstain: Phalloidin Alexa Fluor 633 F-Actin stain; DAPI (blue) nuclear stain at 1:250, 1:50000 for 30-60 min at RT. Magnification: 20X (2X Zoom). (A,C,E,G) - Untreated. (B,D,F,H) - Cells cultured overnight with 50 uM H2O2. (A,B) DAPI (blue) nuclear stain. (C,D) Phalloidin Alex Fluor 633 F-Actin stain. (E,F) Malondialdehyde Antibody. (G,H) Composite. Courtesy of: Dr. Robert Burke, University of Victoria.

ADDITIONAL INFORMATION

Product Categories/Family for anti-MDA antibody	Antibodies; Cancer; Oxidative Stress; Lipid peroxidation; Neuroscience; Neurodegeneration; Alzheimer's Disease
---	--

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/malondialdehyde-antibody-clone-11e3-immunofluorescence-immunocytochemistry-western-blot-elisa/104084>