Mouse Methylglyoxal Monoclonal Antibody | anti-MG antibody

Methylglyoxal Antibody, Clone 9F11: RPE

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA104102	**Mouse**	**Species Independent**	**ELISA, Flow Cytometry, Functional Assay, Immunofluorescence, Immunocytochemistry, Western Blot**


PRODUCT SYNONYMS

Methylglyoxal; Monoclonal Antibody; Methylglyoxal Antibody; Clone 9F11: RPE; Mouse Anti-Methylglyoxal (MG) Monoclonal IgG1; Methylglyoxal Antibody: RPE; 2-Oxopropanal Antibody; 2 oxo propanal Antibody; MG Antibody; Pyruvaldehyde Antibody; Methylglyoxal (MG) Antibody; MG-modified protein Antibody; anti-MG antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Species Independent
Clonality	Monoclonal
Isotype	IgG1
Clone Number	9F11
Specificity	Specific for Methylglyoxal modified proteins. Does not detect free Methylglyoxal. Does not cross-react with Acrolein, Hexanoyl Lysine, Malondialdeyhde, 4-Hydroxy-2-hexenal, 4-Hydroxy nonenal, or Crotonaldehyde modified proteins.
Purity/Purification	Protein G Purified
Concentration	1 mg/ml (varies by lot)
Applicable Applications for anti-MG antibody	ELISA, FCM/FACS (Flow Cytometry), IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Target	Methylglyoxal
Immunogen	Synthetic Methylglyoxal modified Keyhole Limpet Hemocyanin (KLH).
Conjugate	RPE
Storage Buffer	PBS pH 7.4, 50% glycerol, 0.09% Sodium Azide
Preparation and Storage	Store at -20 degree C.


IMAGES



FCM/FACS (Flow Cytometry)
Flow Cytometry analysis using Mouse Anti-Methylglyoxal Monoclonal Antibody, Clone 9F11. Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 90% Methanol. Primary Antibody: Mouse Anti-Methylglyoxal Monoclonal Antibody at 1:50 for 30 min on ice. Secondary Antibody: Goat Anti-Mouse: PE at 1:100 for 20 min at RT. Isotype Control: Non Specific IgG. Cells were subject to oxidative stress by treating with 250 uM H2O2 for 24 hours.



WB (Western Blot)
Western Blot analysis of Methylglyoxal-BSA Conjugate showing detection of 67 kDa Methylglyoxal protein using Mouse Anti-Methylglyoxal Monoclonal Antibody, Clone 9F11. Lane 1: Molecular Weight Ladder (MW). Lane 2: Methylglyoxal-BSA. Lane 3: BSA. Load: 0.5 ug. Block: 5% Skim Milk in TBST. Primary Antibody: Mouse Anti-Methylglyoxal Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:1000 for 60 min at RT. Color Development: ECL solution for 5 min in RT. Predicted/Observed Size: 67 kDa.



ICC (Immunocytochemistry)
Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Methylglyoxal Monoclonal Antibody, Clone 9F11. Tissue: Embryonic kidney epithelial cell line (HEK293). Species: Human. Fixation: 5% Formaldehyde for 5 min. Primary Antibody: Mouse Anti-Methylglyoxal Monoclonal Antibody at 1:50 for 30-60 min at RT. Secondary Antibody: Goat Anti-Mouse Alexa Fluor 488 at 1:1500 for 30-60 min at RT. Counterstain: Phalloidin Alexa Fluor 633 F-Actin stain; DAPI (blue) nuclear stain at 1:250, 1:50000 for 30-60 min at RT. Magnification: 20X (2X Zoom). (A,C,E,G) - Untreated. (B,D,F,H) - Cells cultured overnight with 50 uM H2O2. (A,B) DAPI (blue) nuclear stain. (C,D) Phalloidin Alex Fluor 633 F-Actin stain. (E,F) Methylglyoxal Antibody. (G,H) Composite. Courtesy of: Dr. Robert Burke, University of Victoria.

ADDITIONAL INFORMATION

Product Categories/Family for anti-MG antibody	Antibodies; Cancer; Oxidative Stress; Lipid peroxidation

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: https://stage.aaabiotech.com/monoclonal/methylglyoxal-antibody-clone-9f11-flow-cytometry-immunofluorescence-immunocytochemistry-western-blot-elisa/104102