Mouse Dityrosine Monoclonal Antibody | anti-DT antibody

Dityrosine Antibody, Clone 10A6: FITC

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA104117	**Mouse**	**Species Independent**	**ELISA, Flow Cytometry, Functional Assay, Immunofluorescence, Immunocytochemistry, Western Blot**

PRODUCT SYNONYMS

Dityrosine; Monoclonal Antibody; Dityrosine Antibody; Clone 10A6: FITC; Mouse Anti-Dityrosine (DT) Monoclonal IgG1; Dityrosine Antibody: FITC; Dityrosine (DT) Antibody; DT Antibody; DY Antibody; anti-DT antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Species Independent
Clonality	Monoclonal
Isotype	IgG1
Clone Number	10A6
Specificity	Specific for dityrosine modified proteins. Does not cross-react with 3,5-dibromotyrosine or bromotyrosine modified proteins.
Purity/Purification	Protein G Purified
Concentration	1 mg/ml (varies by lot)
Applicable Applications for anti-DT antibody	ELISA, FCM/FACS (Flow Cytometry), IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Target	Dityrosine
Immunogen	Synthetic Dityrosine conjugated to Keyhole Limpet Hemocyanin (KLH).
Conjugate	FITC
Storage Buffer	PBS pH 7.4, 50% glycerol, 0.09% Sodium Azide
Preparation and Storage	Store at -20 degree C.


IMAGES

WB (Western Blot)
Western Blot analysis of Human Cervical cancer cell line (HeLa) lysate showing detection of Dityrosine protein using Mouse Anti-Dityrosine Monoclonal Antibody, Clone 10A6. Lane 1: Molecular Weight Ladder (MW). Lane 2: HeLa cell lysate. Lane 3: H2O2 treated HeLa cell lysate. Load: 12 ug. Block: 5% Skim Milk in TBST. Primary Antibody: Mouse Anti-Dityrosine Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:2000 for 60 min at RT. Color Development: ECL solution for 5 min in RT.

FCM/FACS (Flow Cytometry)
Flow Cytometry analysis using Mouse Anti-Dityrosine Monoclonal Antibody, Clone 10A6. Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 90% Methanol. Primary Antibody: Mouse Anti-Dityrosine Monoclonal Antibody at 1:50 for 30 min on ice. Secondary Antibody: Goat Anti-Mouse: PE at 1:100 for 20 min at RT. Isotype Control: Non Specific IgG. Cells were subject to oxidative stress by treating with 250 uM H2O2 for 24 hours.



WB (Western Blot)
Western Blot analysis of Dityrosine-BSA Conjugate showing detection of 67 kDa Dityrosine protein using Mouse Anti-Dityrosine Monoclonal Antibody, Clone 10A6. Lane 1: Molecular Weight Ladder (MW). Lane 2: BSA. Lane 3: 3,5-Dibromotyrosine-BSA. Lane 4: Dityrosine-BSA. Lane 5: Bromotyrosine-BSA. Lane 6: 7-ketocholesterol-BSA. Load: 1 ug. Block: 5% Skim Milk in TBST. Primary Antibody: Mouse Anti-Dityrosine Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:2000 for 60 min at RT. Color Development: ECL solution for 5 min in RT. Predicted/Observed Size: 67 kDa.



ICC (Immunocytochemistry)
Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Dityrosine Monoclonal Antibody, Clone 10A6. Tissue: Embryonic kidney epithelial cell line (HEK293). Species: Human. Fixation: 5% Formaldehyde for 5 min. Primary Antibody: Mouse Anti-Dityrosine Monoclonal Antibody at 1:50 for 30-60 min at RT. Secondary Antibody: Goat Anti-Mouse Alexa Fluor 488 at 1:1500 for 30-60 min at RT. Counterstain: Phalloidin Alexa Fluor 633 F-Actin stain; DAPI (blue) nuclear stain at 1:250, 1:50000 for 30-60 min at RT. Localization: Cytoplasmic. Magnification: 20X (2X Zoom). (A,C,E,G) - Untreated. (B,D,F,H) - Cells cultured overnight with 50 uM H2O2. (A,B) DAPI (blue) nuclear stain. (C,D) Phalloidin Alex Fluor 633 F-Actin stain. (E,F) Dityrosine Antibody. (G,H) Composite. Courtesy of: Dr. Robert Burke, University of Victoria.

ADDITIONAL INFORMATION

Product Categories/Family for anti-DT antibody	Antibodies; Cancer; Oxidative Stress; Protein Oxidation; Cell Signaling; Post-translational Modifications; Oxidation

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: https://stage.aaabiotech.com/monoclonal/dityrosine-antibody-clone-10a6-flow-cytometry-immunofluorescence-immunocytochemistry-western-blot-elisa/104117