

Rabbit Peroxiredoxin 3 Polyclonal Antibody | anti-PRDX3 antibody

Anti-Peroxiredoxin 3 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA104271	Rabbit	Human, Mouse, Rat, Monkey	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

Peroxiredoxin 3; Polyclonal Antibody; Anti-Peroxiredoxin 3 Antibody; AOP1; Thioredoxin-dependent peroxide reductase; mitochondrial; Antioxidant protein 1; AOP-1; HBC189; Peroxiredoxin III; Prx-III; Peroxiredoxin-3; Protein MER5 homolog; anti-PRDX3 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat, Monkey
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Peroxiredoxin 3 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	256
Applicable Applications for anti-PRDX3 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Peroxiredoxin 3. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Peroxiredoxin 3 staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Peroxiredoxin 3 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Peroxiredoxin 3 expression in HeLa (A), MCF7 (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-PRDX3 antibody	Rabbit polyclonal antibody to Peroxiredoxin 3
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	5802974
NCBI GeneID	10935
NCBI Accession #	NP_006784.1
NCBI GenBank Nucleotide #	NM_006793.4
UniProt Accession #	P30048; P35690; Q0D2H1; Q13776; Q5T5V2; Q96HK4; B2R7Z0; D3DRC9; E9PH29
Molecular Weight	25,839 Da
NCBI Official Full Name	thioredoxin-dependent peroxide reductase, mitochondrial isoform a
NCBI Official Synonym Full Names	peroxiredoxin 3
NCBI Official Symbol	PRDX3
NCBI Official Synonym Symbols	AOP1; MER5; AOP-1; SP-22; HBC189; PRO1748; prx-III
NCBI Protein Information	thioredoxin-dependent peroxide reductase, mitochondrial; peroxiredoxin-3; peroxiredoxin III; protein MER5 homolog; antioxidant protein 1
UniProt Protein Name	Thioredoxin-dependent peroxide reductase, mitochondrial
UniProt Gene Name	PRDX3
UniProt Synonym Gene Names	AOP1; AOP-1; Prx-III
UniProt Entry Name	PRDX3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/peroxiredoxin-3-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/104271>