

Rabbit anti-Human TRP2 Polyclonal Antibody | anti-DCT antibody

Anti-TRP2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA104469	Rabbit	Human	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

TRP2; Polyclonal Antibody; Anti-TRP2 Antibody; TYRP2; L-dopachrome tautomerase; DCT; DT; L-dopachrome Delta-isomerase; Tyrosinase-related protein 2; TRP-2; anti-DCT antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of TRP2 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	552
Applicable Applications for anti-DCT antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human TRP2. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of TRP2 staining in A375 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of TRP2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of TRP2 expression in A375 (A), Y79 (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-DCT antibody	Rabbit polyclonal antibody to TRP2
---	------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	193788640
NCBI GeneID	1638
NCBI Accession #	NP_001123361.1
NCBI GenBank Nucleotide #	NM_001129889.1
UniProt Accession #	P40126; Q09GT4
Molecular Weight	62,929 Da
NCBI Official Full Name	L-dopachrome tautomerase isoform 2
NCBI Official Synonym Full Names	dopachrome tautomerase
NCBI Official Symbol	DCT
NCBI Official Synonym Symbols	TRP-2; TYRP2
NCBI Protein Information	L-dopachrome tautomerase; DT; TRP2; dopachrome delta-isomerase; tyrosine-related protein 2; L-dopachrome Delta-isomerase; tyrosinase-related protein 2; dopachrome tautomerase (dopachrome delta-isomerase, tyrosine-related protein 2)
UniProt Protein Name	L-dopachrome tautomerase
UniProt Gene Name	DCT
UniProt Synonym Gene Names	TYRP2; DCT; DT; TRP-2; TRP2
UniProt Entry Name	TYRP2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/trp2-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/104469>