

Rabbit NR2F6 Polyclonal Antibody | anti-NR2F6 antibody

Anti-NR2F6 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA104525	Rabbit	Human, Mouse, Rat, Bovine, Dog, Pig	Immunoprecipitation, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

NR2F6; Polyclonal Antibody; Anti-NR2F6 Antibody; EAR2; ERBAL2; Nuclear receptor subfamily 2 group F member 6; V-erbA-related protein 2; EAR-2; anti-NR2F6 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Dog, Pig
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of NR2F6 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	404
Applicable Applications for anti-NR2F6 antibody	IP (Immunoprecipitation), ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human NR2F6. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of TEL staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of TEL staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of TEL expression in CCRFCM (A), Jurkat (B), K562 (C) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-NR2F6 antibody	Rabbit polyclonal antibody to NR2F6
---	-------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	20070199
NCBI GeneID	2063
NCBI Accession #	NP_005225.2
NCBI GenBank Nucleotide #	NM_005234.3
UniProt Accession #	P10588; Q5XGA0; Q6P586; Q9BUE8; B2RC68
Molecular Weight	42,979 Da
NCBI Official Full Name	nuclear receptor subfamily 2 group F member 6
NCBI Official Synonym Full Names	nuclear receptor subfamily 2, group F, member 6
NCBI Official Symbol	NR2F6
NCBI Official Synonym Symbols	EAR2; EAR-2; ERBAL2
NCBI Protein Information	nuclear receptor subfamily 2 group F member 6; ERBA-related gene-2; V-erbA-related protein 2; nuclear receptor V-erbA-related; v-erb-a avian erythroblastic leukemia viral oncogene homolog-like 2
UniProt Protein Name	Nuclear receptor subfamily 2 group F member 6
UniProt Gene Name	NR2F6
UniProt Synonym Gene Names	EAR2; ERBAL2; EAR-2
UniProt Entry Name	NR2F6_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/nr2f6-antibody-immunoprecipitation-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/104525>