

Involucrin Polyclonal Antibody | anti-IVL antibody

Anti-Involucrin Antibody

CATALOG # AAA104565	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot
-------------------------------	--	--

PRODUCT SYNONYMS

Involucrin; Polyclonal Antibody; Anti-Involucrin Antibody; anti-IVL antibody

PRODUCT SPECIFICATIONS


Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Involucrin protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	585
Applicable Applications for anti-IVL antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Source	Rabbit
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Involucrin. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Involucrin staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Involucrin staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Involucrin expression in MCF7 (A), A375 (B), A431 (C), NIH3T3 (D), HT1080 (E), mouse brain (F), rat brain (G) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-IVL antibody	Rabbit polyclonal antibody to Involucrin
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	44890059
NCBI GeneID	3713
NCBI Accession #	NP_005538.2
NCBI GenBank Nucleotide #	NM_005547.2
UniProt Accession #	P07476; Q5T7P4
NCBI Official Full Name	involucrin
NCBI Official Synonym Full Names	involucrin
NCBI Official Symbol	IVL
NCBI Protein Information	involucrin
UniProt Protein Name	Involucrin
UniProt Gene Name	IVL
UniProt Entry Name	INVO_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/involucrin-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/104565>