

Rabbit Desmin Polyclonal Antibody | anti-DES antibody

Anti-Desmin Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA104568	Rabbit	Human, Rat, Bovine, Chicken, Dog, Pig	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

Desmin; Polyclonal Antibody; Anti-Desmin Antibody; anti-DES antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Rat, Bovine, Chicken, Dog, Pig
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Desmin protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	470
Applicable Applications for anti-DES antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Desmin. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES



IF (Immunofluorescence)

Immunofluorescent analysis of Desmin staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Desmin staining in human muscle formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Desmin expression in Sol8 (A), HeLa (B), SJRH30 (C), PC12 (D) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-DES antibody	Rabbit polyclonal antibody to Desmin
---	--------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	55749932
NCBI GeneID	1674
NCBI Accession #	NP_001918.3
NCBI GenBank Nucleotide #	NM_001927.3
UniProt Accession #	P17661; Q15787; Q549R7; Q549R8; Q549R9; Q8IZR1; Q8IZR6; Q8NES2; Q8NEU6; Q8TAC4; Q8TCX2; Q8TD99
Molecular Weight	53,536 Da
NCBI Official Full Name	desmin
NCBI Official Synonym Full Names	desmin
NCBI Official Symbol	DES
NCBI Official Synonym Symbols	CSM1; CSM2; LGMD2R
NCBI Protein Information	desmin; mutant desmin p.K241E; intermediate filament protein
UniProt Protein Name	Desmin
UniProt Gene Name	DES
UniProt Entry Name	DESM_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/desmin-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/104568>