

Rabbit anti-Human Kallikrein 3 Polyclonal Antibody | anti-KLK3 antibody

Anti-Kallikrein 3 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA104629	Rabbit	Human	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

Kallikrein 3; Polyclonal Antibody; Anti-Kallikrein 3 Antibody; APS; Prostate-specific antigen; PSA; Gamma-seminoprotein; Semin; Kallikrein-3; P-30 antigen; Semenogelase; anti-KLK3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Kallikrein 3 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	238
Applicable Applications for anti-KLK3 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Kallikrein 3. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Kallikrein 3 staining in LNCaP cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Kallikrein 3 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Kallikrein 3 expression in LNCaP (A) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-KLK3 antibody	Rabbit polyclonal antibody to Kallikrein 3
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	71834853
NCBI GeneID	354
NCBI Accession #	NP_001025218.1
NCBI GenBank Nucleotide #	NM_001030047.1
UniProt Accession #	P07288; Q15096; Q16272; Q86TG8; Q8IXI4; C9JXH3; G3V0H4; G3XAE3
Molecular Weight	25,116 Da
NCBI Official Full Name	prostate-specific antigen isoform 3 preproprotein
NCBI Official Synonym Full Names	kallikrein-related peptidase 3
NCBI Official Symbol	KLK3
NCBI Official Synonym Symbols	APS; PSA; hK3; KLK2A1
NCBI Protein Information	prostate-specific antigen; seminin; P-30 antigen; kallikrein-3; semenogelase; gamma-seminoprotein; prostate specific antigen
UniProt Protein Name	Prostate-specific antigen
UniProt Gene Name	KLK3
UniProt Synonym Gene Names	APS; PSA; Semin
UniProt Entry Name	KLK3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/kallikrein-3-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/104629>