

Rabbit Creatine Kinase M Polyclonal Antibody | anti-CKM antibody

Anti-Creatine Kinase M Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA104638	Rabbit	Human, Mouse, Rat, Monkey	Immunoprecipitation, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Creatine Kinase M; Polyclonal Antibody; Anti-Creatine Kinase M Antibody; CKMM; Creatine kinase M-type; Creatine kinase M chain; M-CK; anti-CKM antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Monkey
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Creatine Kinase M protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	381
Applicable Applications for anti-CKM antibody	IP (Immunoprecipitation), ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human Creatine Kinase M. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Creatine Kinase M staining in NIH3T3 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Creatine Kinase M staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Creatine Kinase M expression in HeLa (A), NIH3T3 (B), H9C2 (C) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-CKM antibody	Rabbit polyclonal antibody to Creatine Kinase M
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	21536288
NCBI GeneID	1158
NCBI Accession #	NP_001815.2
NCBI GenBank Nucleotide #	NM_001824.4
UniProt Accession #	P06732; Q96QL9
Molecular Weight	43,101 Da
NCBI Official Full Name	creatine kinase M-type
NCBI Official Synonym Full Names	creatine kinase, muscle
NCBI Official Symbol	CKM
NCBI Official Synonym Symbols	CKMM; M-CK
NCBI Protein Information	creatine kinase M-type; creatine kinase M chain
UniProt Protein Name	Creatine kinase M-type
UniProt Gene Name	CKM
UniProt Synonym Gene Names	CKMM
UniProt Entry Name	KCRM_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/creatine-kinase-m-antibody-immunoprecipitation-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/104638>