

Rabbit anti-Human, Pig CHK2 Polyclonal Antibody | anti-CHEK2 antibody

Anti-CHK2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA104756	Rabbit	Human, Pig	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CHK2; Polyclonal Antibody; Anti-CHK2 Antibody; CDS1; RAD53; Serine/threonine-protein kinase Chk2; CHK2 checkpoint homolog; Cds1 homolog; Hucds1; hCds1; Checkpoint kinase 2; anti-CHEK2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Pig
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of CHK2 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	586
Applicable Applications for anti-CHEK2 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human CHK2. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of CHK2 staining in HL60 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of CHK2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of CHK2 expression in Jurkat (A), HL60 (B), HT29 (C) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-CHEK2 antibody

Rabbit polyclonal antibody to CHK2

NCBI AND UNIPROT INFORMATION

NCBI GI #	54112407
NCBI GeneID	11200
NCBI Accession #	NP_001005735.1
NCBI GenBank Nucleotide #	NM_001005735.1
UniProt Accession #	O96017; Q6QA03; Q6QA04; Q6QA05; Q6QA06; Q6QA07; Q6QA08; Q6QA10; A8K3Y9; B7ZBF3; B7ZBF4; B7ZBF5
Molecular Weight	36,157 Da
NCBI Official Full Name	serine/threonine-protein kinase Chk2 isoform c
NCBI Official Synonym Full Names	checkpoint kinase 2
NCBI Official Symbol	CHEK2
NCBI Official Synonym Symbols	CDS1; CHK2; LFS2; RAD53; hCds1; HuCds1; PP1425
NCBI Protein Information	serine/threonine-protein kinase Chk2; cds1 homolog; CHK2 checkpoint homolog; checkpoint-like protein CHK2
UniProt Protein Name	Serine/threonine-protein kinase Chk2
UniProt Gene Name	CHEK2
UniProt Synonym Gene Names	CDS1; CHK2; RAD53; Hucds1; hCds1
UniProt Entry Name	CHK2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/chk2-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/104756>