

Rabbit anti-Human Cytokeratin 16 Polyclonal Antibody | anti-KRT16 antibody

Anti-Cytokeratin 16 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA104813	Rabbit	Human	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Cytokeratin 16; Polyclonal Antibody; Anti-Cytokeratin 16 Antibody; KRT16A; Keratin; type I cytoskeletal 16; Cytokeratin-16; CK-16; Keratin-16; K16; anti-KRT16 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Cytokeratin 16 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	473
Applicable Applications for anti-KRT16 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Cytokeratin 16. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Cytokeratin 16 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Cytokeratin 16 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)
Western blot analysis of Cytokeratin 16 expression in HeLa (A) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-KRT16 antibody	Rabbit polyclonal antibody to Cytokeratin 16
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	24430192
NCBI GeneID	3868
NCBI Accession #	NP_005548.2
NCBI GenBank Nucleotide #	NM_005557.3
UniProt Accession #	P08779; P30654; Q16402; Q9UBG8; A8K488
Molecular Weight	51,268 Da
NCBI Official Full Name	keratin, type I cytoskeletal 16
NCBI Official Synonym Full Names	keratin 16
NCBI Official Symbol	KRT16
NCBI Official Synonym Symbols	K16; PC1; CK16; K1CP; NEPPK; FNEPPK; KRT16A
NCBI Protein Information	keratin, type I cytoskeletal 16; CK-16; keratin-16; cytokeratin 16; cytokeratin-16; focal non-epidermolytic palmoplantar keratoderma
UniProt Protein Name	Keratin, type I cytoskeletal 16
UniProt Gene Name	KRT16
UniProt Synonym Gene Names	KRT16A; CK-16; K16
UniProt Entry Name	K1C16_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.