

Rabbit CDC37 Polyclonal Antibody | anti-CDC37 antibody

Anti-CDC37 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA104969	Rabbit	Human, Mouse, Rat, Monkey	Immunoprecipitation, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CDC37; Polyclonal Antibody; Anti-CDC37 Antibody; CDC37A; Hsp90 co-chaperone Cdc37; Hsp90 chaperone protein kinase-targeting subunit; p50Cdc37; anti-CDC37 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Monkey
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of CDC37 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	378
Applicable Applications for anti-CDC37 antibody	IP (Immunoprecipitation), ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CDC37. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of CDC37 staining in K562 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of CDC37 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of CDC37 expression in Jurkat (A), K562 (B), HL60 (C), NIH3T3 (D) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-CDC37 antibody	Rabbit polyclonal antibody to CDC37
---	-------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	5901922
NCBI GeneID	11140
NCBI Accession #	NP_008996.1
NCBI GenBank Nucleotide #	NM_007065.3
UniProt Accession #	Q16543; Q53YA2
Molecular Weight	44,468 Da
NCBI Official Full Name	hsp90 co-chaperone Cdc37
NCBI Official Synonym Full Names	cell division cycle 37
NCBI Official Symbol	CDC37
NCBI Official Synonym Symbols	P50CDC37
NCBI Protein Information	hsp90 co-chaperone Cdc37; cell division cycle 37 homolog; CDC37 cell division cycle 37 homolog; hsp90 chaperone protein kinase-targeting subunit; CDC37 (cell division cycle 37, S. cerevisiae, homolog)
UniProt Protein Name	Hsp90 co-chaperone Cdc37
UniProt Gene Name	CDC37
UniProt Synonym Gene Names	CDC37A
UniProt Entry Name	CDC37_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/polyclonal/cdc37-antibody-immunoprecipitation-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/104969>