

Rabbit Histone H2A.X Polyclonal Antibody | anti-H2AFX antibody

Anti-Histone H2A.X Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA105031	Rabbit	Human, Rabbit	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

Histone H2A.X; Polyclonal Antibody; Anti-Histone H2A.X Antibody; H2AX; Histone H2AX; H2a/x; anti-H2AFX antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Rabbit
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Histone H2A.X protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	143
Applicable Applications for anti-H2AFX antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Histone H2A.X. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Histone H2A.X staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of Histone H2A.X staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Histone H2A.X expression in Hela (A), MCF7 (B), Jurkat (C) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-H2AFX antibody	Rabbit polyclonal antibody to Histone H2A.X
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504253
NCBI GeneID	3014
NCBI Accession #	NP_002096.1
NCBI GenBank Nucleotide #	NM_002105.2
UniProt Accession #	P16104; Q4ZGJ7; Q6IAS5
Molecular Weight	15,145 Da
NCBI Official Full Name	histone H2AX
NCBI Official Synonym Full Names	H2A histone family, member X
NCBI Official Symbol	H2AFX
NCBI Official Synonym Symbols	H2AX; H2A.X; H2A/X
NCBI Protein Information	histone H2AX; H2AX histone; histone H2A.x
UniProt Protein Name	Histone H2AX
UniProt Gene Name	H2AFX
UniProt Synonym Gene Names	H2AX; H2a/x
UniProt Entry Name	H2AX_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/histone-h2ax-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/105031>