

Rabbit anti-Human Cystatin B Polyclonal Antibody | anti-CSTB antibody

Anti-Cystatin B Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA105045	Rabbit	Human	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

Cystatin B; Polyclonal Antibody; Anti-Cystatin B Antibody; CST6; STFB; Cystatin-B; CPI-B; Liver thiol proteinase inhibitor; Stefin-B; anti-CSTB antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Cystatin B protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	98
Applicable Applications for anti-CSTB antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Cystatin B. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Cystatin B staining in EOC20 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of Cystatin B staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Cystatin B expression in JAR (A), EOC20 (B), U87MG (C) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-CSTB antibody	Rabbit polyclonal antibody to Cystatin B
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503117
NCBI GeneID	1476
NCBI Accession #	NP_000091.1
NCBI GenBank Nucleotide #	NM_000100.3
UniProt Accession #	P04080
Molecular Weight	11,140 Da
NCBI Official Full Name	cystatin-B
NCBI Official Synonym Full Names	cystatin B (stefin B)
NCBI Official Symbol	CSTB
NCBI Official Synonym Symbols	PME; ULD; CST6; EPM1; STFB; EPM1A
NCBI Protein Information	cystatin-B; CPI-B; liver thiol proteinase inhibitor
UniProt Protein Name	Cystatin-B
UniProt Gene Name	CSTB
UniProt Synonym Gene Names	CST6; STFB
UniProt Entry Name	CYTB_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cystatin-b-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/105045>