

Rabbit anti-Human, Mouse DNA-PKcs Polyclonal Antibody | anti-PRKDC antibody

Anti-DNA-PKcs Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA105074	Rabbit	Human, Mouse	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

DNA-PKcs; Polyclonal Antibody; Anti-DNA-PKcs Antibody; HYRC; HYRC1; DNA-dependent protein kinase catalytic subunit; DNA-PK catalytic subunit; DNPk1; p460; anti-PRKDC antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of DNA-PKcs protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	4097
Applicable Applications for anti-PRKDC antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human DNA-PKcs. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of DNA-PKcs staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of DNA-PKcs staining in human tonsil formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of DNA-PKcs expression in HeLa (A), MCF7 (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-PRKDC antibody

Rabbit polyclonal antibody to DNA-PKcs

NCBI AND UNIPROT INFORMATION

NCBI GI #	126032350
NCBI GeneID	5591
NCBI Accession #	NP_001075109.1
NCBI GenBank Nucleotide #	NM_001081640.1
UniProt Accession #	P78527; P78528; Q13327; Q13337; Q14175; Q59H99; Q7Z611; Q96SE6; Q9UME3
Molecular Weight	465,501 Da
NCBI Official Full Name	DNA-dependent protein kinase catalytic subunit isoform 2
NCBI Official Synonym Full Names	protein kinase, DNA-activated, catalytic polypeptide
NCBI Official Symbol	PRKDC
NCBI Official Synonym Symbols	HYRC; p350; DNAPK; DNP1K1; HYRC1; IMD26; XRCC7; DNA-PKcs
NCBI Protein Information	DNA-dependent protein kinase catalytic subunit; p460; DNA-PK catalytic subunit; hyper-radiosensitivity of murine scid mutation, complementing 1
UniProt Protein Name	DNA-dependent protein kinase catalytic subunit
UniProt Gene Name	PRKDC
UniProt Synonym Gene Names	HYRC; HYRC1; DNA-PK catalytic subunit; DNA-PKcs
UniProt Entry Name	PRKDC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/dna-pkcs-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/105074>