

Rabbit Cytokeratin 17 Polyclonal Antibody | anti-KRT17 antibody

Anti-Cytokeratin 17 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA105138	Rabbit	Human, Rat, Dog, Rabbit	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Cytokeratin 17; Polyclonal Antibody; Anti-Cytokeratin 17 Antibody; Keratin; type I cytoskeletal 17; 39.1; Cytokeratin-17; CK-17; Keratin-17; K17; anti-KRT17 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Rat, Dog, Rabbit
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Cytokeratin 17 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	432
Applicable Applications for anti-KRT17 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Cytokeratin 17. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Cytokeratin 17 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Cytokeratin 17 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Cytokeratin 17 expression in PC12 (A), HeLa (B), A431 (C), NIH3T3 (D) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-KRT17 antibody

Rabbit polyclonal antibody to Cytokeratin 17

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557701
NCBI GeneID	3872
NCBI Accession #	NP_000413.1
NCBI GenBank Nucleotide #	NM_000422.2
UniProt Accession #	Q04695; Q6IP98; Q8N1P6; A5Z1M9; A5Z1N0; A5Z1N1; A5Z1N2; A6NDV6; A6NKQ2
Molecular Weight	48,106 Da
NCBI Official Full Name	keratin, type I cytoskeletal 17
NCBI Official Synonym Full Names	keratin 17
NCBI Official Symbol	KRT17
NCBI Official Synonym Symbols	PC; K17; PC2; PCHC1
NCBI Protein Information	keratin, type I cytoskeletal 17; 39.1; CK-17; keratin-17; cytokeratin-17; keratin 17 epitope S1; keratin 17 epitope S2; keratin 17 epitope S4
UniProt Protein Name	Keratin, type I cytoskeletal 17
UniProt Gene Name	KRT17
UniProt Synonym Gene Names	CK-17; K17
UniProt Entry Name	K1C17_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/cytokeratin-17-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/105138>