

Rabbit ACTN2 Polyclonal Antibody | anti-ACTN2 antibody

Anti-ACTN2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA105146	Rabbit	Human, Mouse, Rat, Bovine, Chicken	Immunoprecipitation, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

ACTN2; Polyclonal Antibody; Anti-ACTN2 Antibody; Alpha-actinin-2; Alpha-actinin skeletal muscle isoform 2; F-actin cross-linking protein; ACTN3; Alpha-actinin-3; Alpha-actinin skeletal muscle isoform 3; anti-ACTN2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Chicken
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of ACTN2 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	894
Applicable Applications for anti-ACTN2 antibody	IP (Immunoprecipitation), ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human ACTN2. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of BAX staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of BAX staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of BAX expression in mouse brain (A), rat brain (B), A2780 (C), MCF7 (D) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-ACTN2 antibody

Rabbit polyclonal antibody to ACTN2

NCBI AND UNIPROT INFORMATION

NCBI GI #	4501893
NCBI GeneID	88
NCBI Accession #	NP_001094.1
NCBI GenBank Nucleotide #	NM_001103.3
UniProt Accession #	P35609; Q86TF4; Q86TI8; B1ANE4; B2RCS5
Molecular Weight	103,920 Da
NCBI Official Full Name	alpha-actinin-2 isoform 1
NCBI Official Synonym Full Names	actinin, alpha 2
NCBI Official Symbol	ACTN2
NCBI Official Synonym Symbols	CMD1AA
NCBI Protein Information	alpha-actinin-2; F-actin cross-linking protein; alpha-actinin skeletal muscle
UniProt Protein Name	Alpha-actinin-2
UniProt Gene Name	ACTN2
UniProt Entry Name	ACTN2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/actn2-antibody-immunoprecipitation-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/105146>