

Connexin 43 Polyclonal Antibody | anti-GJA1 antibody

Anti-Connexin 43 Antibody

CATALOG #	REACTIVITY	APPLICATIONS
AAA105156	Human, Mouse, Rat, Bovine, Chicken, Dog, Rabbit	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Connexin 43; Polyclonal Antibody; Anti-Connexin 43 Antibody; GJAL; Gap junction alpha-1 protein; Connexin-43; Cx43; Gap junction 43 kDa heart protein; anti-GJA1 antibody

PRODUCT SPECIFICATIONS


Reactivity	Human, Mouse, Rat, Bovine, Chicken, Dog, Rabbit
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Connexin 43 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	382
Applicable Applications for anti-GJA1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Source	Rabbit
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Connexin 43. The exact sequence is proprietary.
Preparation and Storage	Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Connexin 43 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Connexin 43 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)
 Western blot analysis of Connexin 43 expression in C6 (A), HeLa (B), C2C12 (C), mouse heart (D), rat brain (E) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-GJA1 antibody	Rabbit polyclonal antibody to Connexin 43
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504001
NCBI GeneID	2697
NCBI Accession #	NP_000156.1
NCBI GenBank Nucleotide #	NM_000165.4
UniProt Accession #	P17302; Q6FHU1; Q9Y5I8; B2R5U9
Molecular Weight	43,008 Da
NCBI Official Full Name	gap junction alpha-1 protein
NCBI Official Synonym Full Names	gap junction protein, alpha 1, 43kDa
NCBI Official Symbol	GJA1
NCBI Official Synonym Symbols	HSS; CMDR; CX43; GJAL; ODDD; AVSD3; HLHS1
NCBI Protein Information	gap junction alpha-1 protein; connexin 43; connexin-43; gap junction 43 kDa heart protein
UniProt Protein Name	Gap junction alpha-1 protein
UniProt Gene Name	GJA1
UniProt Synonym Gene Names	GJAL; Cx43
UniProt Entry Name	CXA1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.