

Rabbit Cyclin A1/2 Polyclonal Antibody | anti-CCNA1 antibody

Anti-Cyclin A1/2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA105252	Rabbit	Human, Mouse, Rat, Bovine, Chicken, Rabbit	Chromatin Immunoprecipitation, Immunoprecipitation, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Cyclin A1/2; Polyclonal Antibody; Anti-Cyclin A1/2 Antibody; CCNA1; Cyclin-A1; CCNA2; CCN1; CCNA; Cyclin-A2; Cyclin-A; anti-CCNA1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Chicken, Rabbit
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Cyclin A1/2 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	464
Applicable Applications for anti-CCNA1 antibody	ChIP (Chromatin immunoprecipitation), ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Cyclin A1/2. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of TBX15/18 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of TBX15/18 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of TBX15/18 expression in Jurkat (A), HepG2 (B), LOVO (C) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-CCNA1 antibody

Rabbit polyclonal antibody to Cyclin A1/2

NCBI AND UNIPROT INFORMATION

NCBI GI #	161377468
NCBI GeneID	8900
NCBI Accession #	NP_001104515.1
NCBI GenBank Nucleotide #	NM_001111045.1
UniProt Accession #	P78396; Q5T3V0; Q5U0G2; Q8IY91; B7Z7E3
Molecular Weight	47,494 Da
NCBI Official Full Name	cyclin-A1 isoform b
NCBI Official Synonym Full Names	cyclin A1
NCBI Official Symbol	CCNA1
NCBI Official Synonym Symbols	CT146
NCBI Protein Information	cyclin-A1
UniProt Protein Name	Cyclin-A1
UniProt Gene Name	CCNA1
UniProt Entry Name	CCNA1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/cyclin-a1-2-antibody-chromatin-immunoprecipitation-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/105252>