

Cytokeratin 5 Polyclonal Antibody | anti-KRT5 antibody

Anti-Cytokeratin 5 Antibody

CATALOG # AAA105277	REACTIVITY Human, Mouse, Rat, Bovine, Pig	APPLICATIONS Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot
-------------------------------	---	--


PRODUCT SYNONYMS

Cytokeratin 5; Polyclonal Antibody; Anti-Cytokeratin 5 Antibody; Keratin; type II cytoskeletal 5; 58 kDa cytokeratin; Cytokeratin-5; CK-5; Keratin-5; K5; Type-II keratin Kb5; anti-KRT5 antibody

PRODUCT SPECIFICATIONS


Reactivity	Human, Mouse, Rat, Bovine, Pig
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Cytokeratin 5 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	590
Applicable Applications for anti-KRT5 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Source	Rabbit
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Cytokeratin 5. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Cytokeratin 5 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Cytokeratin 5 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Cytokeratin 5 expression in DU145 (A), HeLa (B), MCF7 (C) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-KRT5 antibody

Rabbit polyclonal antibody to Cytokeratin 5

NCBI AND UNIPROT INFORMATION

NCBI GI #	119395754
NCBI GeneID	3852
NCBI Accession #	NP_000415.2
NCBI GenBank Nucleotide #	NM_000424.3
UniProt Accession #	P13647; Q6PI71; Q6UBJ0; Q8TA91
Molecular Weight	62,378 Da
NCBI Official Full Name	keratin, type II cytoskeletal 5
NCBI Official Synonym Full Names	keratin 5
NCBI Official Symbol	KRT5
NCBI Official Synonym Symbols	K5; CK5; DDD; DDD1; EBS2; KRT5A
NCBI Protein Information	keratin, type II cytoskeletal 5; CK-5; cytokeratin-5; 58 kda cytokeratin; type-II keratin Kb5; epidermolysis bullosa simplex 2 Dowling-Meara/Kobner/Weber-Cockayne types; keratin 5 (epidermolysis bullosa simplex, Dowling-Meara/Kobner/Weber-Cockayne types)
UniProt Protein Name	Keratin, type II cytoskeletal 5
UniProt Gene Name	KRT5
UniProt Synonym Gene Names	CK-5; K5
UniProt Entry Name	K2C5_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/cytokeratin-5-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/105277>