

Rabbit RB1 (pS780) Polyclonal Antibody | anti-RB1 antibody

Anti-RB1 (pS780) Phospho Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA105310	Rabbit	Human, Bovine, Dog, Pig	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

RB1 (pS780); Polyclonal Antibody; Anti-RB1 (pS780) Phospho Antibody; Retinoblastoma-associated protein; p105-Rb; pRb; Rb; pp110; anti-RB1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Bovine, Dog, Pig
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of RB1 (pS780) protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	928
Applicable Applications for anti-RB1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human RB1. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of RB1 (pS780) staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of RB1 (pS780) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of RB1 (pS780) expression in Jurkat FBS-treated (A), HeLa (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-RB1 antibody	Rabbit polyclonal antibody to RB1 (pS780)
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	108773787
NCBI GeneID	5925
NCBI Accession #	NP_000312.2
NCBI GenBank Nucleotide #	NM_000321.2
UniProt Accession #	P06400; P78499; Q5VW46; Q8IZL4; A8K5E3
Molecular Weight	106,159 Da
NCBI Official Full Name	retinoblastoma-associated protein
NCBI Official Synonym Full Names	retinoblastoma 1
NCBI Official Symbol	RB1
NCBI Official Synonym Symbols	RB; pRb; OSRC; pp110; p105-Rb; PPP1R130
NCBI Protein Information	retinoblastoma-associated protein; retinoblastoma susceptibility protein; prepro-retinoblastoma-associated protein; protein phosphatase 1, regulatory subunit 130
UniProt Protein Name	Retinoblastoma-associated protein
UniProt Gene Name	RB1
UniProt Synonym Gene Names	Rb
UniProt Entry Name	RB_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/rb1-ps780-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/105310>