

Rabbit anti-Human, Rat MMP8 Polyclonal Antibody | anti-MMP8 antibody

Anti-MMP8 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA105340	Rabbit	Human, Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MMP8; Polyclonal Antibody; Anti-MMP8 Antibody; CLG1; Neutrophil collagenase; Matrix metalloproteinase-8; MMP-8; PMNL collagenase; PMNL-CL; anti-MMP8 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of MMP8 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	467
Applicable Applications for anti-MMP8 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human MMP8. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of MMP8 staining in SW620 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of MMP8 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of MMP8 expression in SJRH30 (A), PMNL (B), Sw620 (C) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-MMP8 antibody

Rabbit polyclonal antibody to MMP8

NCBI AND UNIPROT INFORMATION

NCBI GI #	4505221
NCBI GeneID	4317
NCBI Accession #	NP_002415.1
NCBI GenBank Nucleotide #	NM_002424.2
UniProt Accession #	P22894; Q45F99
Molecular Weight	53,412 Da
NCBI Official Full Name	neutrophil collagenase preproprotein
NCBI Official Synonym Full Names	matrix metalloproteinase 8 (neutrophil collagenase)
NCBI Official Symbol	MMP8
NCBI Official Synonym Symbols	HNC; CLG1; MMP-8; PMNL-CL
NCBI Protein Information	neutrophil collagenase; PMNL collagenase; matrix metalloproteinase-8; matrix metalloproteinase 8 (neutrophil collagenase)
UniProt Protein Name	Neutrophil collagenase
UniProt Gene Name	MMP8
UniProt Synonym Gene Names	CLG1; MMP-8; PMNL-CL
UniProt Entry Name	MMP8_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/mmp8-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/105340>