

Rabbit IKK gamma Polyclonal Antibody | anti-IKBKG antibody

Anti-IKK gamma Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA105345	Rabbit	Human, Mouse, Rat, Bovine, Rabbit	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

IKK gamma; Polyclonal Antibody; Anti-IKK gamma Antibody; FIP3; NEMO; NF-kappa-B essential modulator; FIP-3; Ikb kinase-associated protein 1; IKKAP1; Inhibitor of nuclear factor kappa-B kinase subunit gamma; I-kappa-B kinase subunit gamma; IKK-gamma; IKKG; Ikb kinase subunit gamma; NF-kappa-B essential modifier; anti-IKBKG antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Rabbit
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of IKK gamma protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	487
Applicable Applications for anti-IKBKG antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human IKK gamma. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of IKK gamma staining in HL60 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of IKK gamma staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of IKK gamma expression in Jurkat (A), HL60 (B), HeLa (C), NIH3T3 (D), Raw264.7 (E), PC12 (F), rat heart (G) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-IKBKG antibody	Rabbit polyclonal antibody to IKK gamma
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	223671870
NCBI GeneID	8517
NCBI Accession #	NP_001093326.2
NCBI GenBank Nucleotide #	NM_001099856.3
UniProt Accession #	Q9Y6K9; Q7LBY6; Q7Z7F1
Molecular Weight	36,953 Da
NCBI Official Full Name	NF-kappa-B essential modulator isoform b
NCBI Official Synonym Full Names	inhibitor of kappa light polypeptide gene enhancer in B-cells, kinase gamma
NCBI Official Symbol	IKBKG
NCBI Official Synonym Symbols	IP; IP1; IP2; FIP3; IPD2; NEMO; FIP-3; Fip3p; IMD33; AMCBX1; ZC2HC9; IKK-gamma
NCBI Protein Information	NF-kappa-B essential modulator; IKKG; IKKAP1; incontinentia pigmenti; Ikb kinase gamma subunit; ikB kinase subunit gamma; NFkappaB essential modulator; NF-kappa-B essential modifier; I-kappa-B kinase subunit gamma; ikB kinase-associated protein 1; inhibitor of nuclear factor kappa-B kinase subunit gamma
UniProt Protein Name	NF-kappa-B essential modulator
UniProt Gene Name	IKBKG
UniProt Synonym Gene Names	FIP3; NEMO; NEMO; IKKAP1; I-kappa-B kinase subunit gamma; IKK-gamma; IKKG; Ikb kinase subunit gamma
UniProt Entry Name	NEMO_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.