

Human CXCL7 ELISA Kit | PPBP elisa kit

Human CXCL7 ELISA Kit

CATALOG # AAA105769	REACTIVITY Human	APPLICATIONS ELISA
-------------------------------	----------------------------	------------------------------


PRODUCT SYNONYMS

CXCL7; Human CXCL7 ELISA Kit; CTAP3; SCYB7; TGB1; THBGB1; Platelet basic protein; PBP; C-X-C motif chemokine 7; Leukocyte-derived growth factor; LDGF; Macrophage-derived growth factor; MDGF; Small-inducible cytokine B7; PPBP elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	The Human CXCL7 ELISA Kit allows for the detection and quantification of endogenous levels of natural and/or recombinant Human CXCL7 proteins within the range of 15.6 pg/ml - 1000 pg/ml.
Sequence Length	128
Samples	Serum, plasma
Sensitivity	Typically less than 7 pg/ml.
Applicable Applications for PPBP elisa kit	ELISA (Sandwich ELISA)
Preparation and Storage	Shipped and store at 4 degree C for 6 months, store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for PPBP elisa kit	Principle of the Assay: Human CXCL7 ELISA (Enzyme-Linked Immunosorbent Assay) kit is an in vitro enzyme-linked immunosorbent assay for the quantitative measurement of Human CXCL7 in Cell Culture Supernatants, Serum, Plasma. This assay employs an antibody specific for Human CXCL7 coated on a 96-well plate. Standards and samples are pipetted into the wells and CXCL7 present in a sample is bound to the wells by the immobilized antibody. The wells are washed and biotinylated anti-Human CXCL7 antibody is added. After washing away unbound biotinylated antibody, HRP-conjugated streptavidin is pipetted to the wells. The wells are again washed, a TMB substrate solution is added to the wells and color develops in proportion to the amount of CXCL7 bound. The Stop Solution changes the color from blue to yellow, and the intensity of the color is measured at 450 nm. Background/Introduction: Chemokine (C-X-C motif) ligand 7 (CXCL7), also called PPBP or SCYB7, is a human gene. The encoded protein, Chemokine (C-X-C motif) ligand is a small cytokine belonging to the CXC chemokine family. CXCL7 is mapped to 4q13.3. It is a protein that is released in large amounts from platelets following their activation. CXCL7 is the precursor of the 2 platelet alpha-granule proteins, platelet basic protein (PBP) and connective tissue-activating peptide III (CTAP3). It stimulates various processes including mitogenesis, synthesis of extracellular matrix, glucose metabolism and synthesis of plasminogen activator. It also stimulates the formation and secretion of plasminogen activator by synovial cells.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4505981
NCBI GeneID	5473
NCBI Accession #	NP_002695.1
NCBI GenBank Nucleotide #	NM_002704.3
UniProt Accession #	P02775; Q6IBJ8; B2R5F3; AAB46877
Molecular Weight	13,894 Da
NCBI Official Full Name	platelet basic protein preproprotein
NCBI Official Synonym Full Names	pro-platelet basic protein (chemokine (C-X-C motif) ligand 7)
NCBI Official Symbol	PPBP
NCBI Official Synonym Symbols	PBP; TC1; TC2; TGB; LDGF; MDGF; TGB1; B-TG1; CTAP3; CXCL7; NAP-2; SCYB7; THBGB; LA-PF4; THBGB1; Beta-TG; CTAPIII; CTAP-III
NCBI Protein Information	platelet basic protein; thrombocidin 1; thrombocidin 2; beta-thromboglobulin; CXC chemokine ligand 7; C-X-C motif chemokine 7; thromboglobulin, beta-1; small inducible cytokine B7; small-inducible cytokine B7; leukocyte-derived growth factor; low-affinity platelet factor IV; neutrophil-activating peptide 2; neutrophil-activating peptide-2; macrophage-derived growth factor; connective tissue-activating peptide III; small inducible cytokine subfamily B, member 7
UniProt Protein Name	Platelet basic protein
UniProt Gene Name	PPBP
UniProt Synonym Gene Names	CTAP3; CXCL7; SCYB7; TGB1; THBGB1; PBP; LDGF; MDGF; CTAP-III; CTAP-III(1-81); Beta-TG; NAP-2(74); NAP-2(73); NAP-2; NAP-2(1-66); NAP-2(1-63)
UniProt Entry Name	CXCL7_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.