

Mouse IL-17 ELISA Kit | IL17A elisa kit

Mouse IL-17 ELISA Kit

CATALOG # AAA106059	REACTIVITY Mouse	APPLICATIONS ELISA
-------------------------------	----------------------------	------------------------------


PRODUCT SYNONYMS

IL-17; Mouse IL-17 ELISA Kit; CTLA8; IL17; Interleukin-17A; IL-17A; Cytotoxic T-lymphocyte-associated antigen 8; CTLA-8; IL17A elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	The Mouse IL-17 ELISA Kit allows for the detection and quantification of endogenous levels of natural and/or recombinant Mouse IL-17 proteins within the range of 7.8 pg/ml - 500 pg/ml.
Sequence Length	158
Samples	Cell Culture Supernatants, Serum, Plasma, Tissue Homogenates
Assay Type	Quantitative Sandwich
Sensitivity	<4 pg/ml
Applicable Applications for IL17A elisa kit	ELISA (Sandwich ELISA)
Preparation and Storage	Shipped and store at 4 degree C for 6 months, store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for IL17A elisa kit	Principle of the Assay: The Mouse IL-17 ELISA (Enzyme-Linked Immunosorbent Assay) kit is an in vitro enzyme-linked immunosorbent assay for the quantitative measurement of Mouse IL-17 in Cell Culture Supernatants, Serum, Plasma, Tissue Homogenates. This assay employs an antibody specific for Mouse IL-17 coated on a 96-well plate. Standards and samples are pipetted into the wells and IL-17 present in a sample is bound to the wells by the immobilized antibody. The wells are washed and biotinylated anti-Mouse IL-17 antibody is added. After washing away unbound biotinylated antibody, HRP-conjugated streptavidin is pipetted to the wells. The wells are again washed, a TMB substrate solution is added to the wells and color develops in proportion to the amount of IL-17 bound. The Stop Solution changes the color from blue to yellow, and the intensity of the color is measured at 450 nm.!!Background/Introduction: IL-17 is an inflammatory cytokine produced primarily by a unique lineage of CD4 T cells that plays critical roles in the pathogenesis of multiple autoimmune diseases. Interleukin-17 is expressed by activated T cells and is 57% identical to the 17- to 26-kD secretory glycoprotein encoded by gene 13 of the herpesvirus saimiri (HVS-13). IL17 induces nuclear factor kappa-B and the expression of IL6, intercellular adhesion molecule-1, granulocyte macrophage colony-stimulating factor,and prostaglandin E2, as well as the maturation of CD34 positive hematopoietic precursors into neutrophils. Anti-IL17 antibodies significantly inhibited osteoclast formation induced by culture media of RA synovial tissues.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	6754324
NCBI GeneID	16171
NCBI Accession #	NP_034682.1
NCBI GenBank Nucleotide #	NM_010552.3
UniProt Accession #	Q62386; Q60971
Molecular Weight	17,490 Da
NCBI Official Full Name	interleukin-17A
NCBI Official Synonym Full Names	interleukin 17A
NCBI Official Symbol	Il17a
NCBI Official Synonym Symbols	Il17; Ctl8; IL-17; Ctl8; IL-17A
NCBI Protein Information	interleukin-17A; cytotoxic T-lymphocyte-associated antigen 8
UniProt Protein Name	Interleukin-17A
UniProt Gene Name	Il17a
UniProt Synonym Gene Names	Ctl8; Il17; IL-17; IL-17A; CTLA-8
UniProt Entry Name	IL17_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/il17a-mouse-elisa-kits/il-17/106059>