

Rabbit PKC zeta Polyclonal Antibody | anti-PRKCZ antibody

Anti-PKC zeta Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA106138	Rabbit	Human, Mouse, Rat, Monkey, Rabbit, Zebrafish	Immunoprecipitation, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

PKC zeta; Polyclonal Antibody; Anti-PKC zeta Antibody; PKC2; Protein kinase C zeta type; nPKC-zeta; anti-PRKCZ antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Monkey, Rabbit, Zebrafish
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of PKC zeta protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	409
Applicable Applications for anti-PRKCZ antibody	IP (Immunoprecipitation), ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human PKC zeta. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of NF-kappaB p65 (AcK310) staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of NF-kappaB p65 (AcK310) staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)
 Western blot analysis of NF-kappaB p65 (AcK310) expression in HeLa (A), mouse lung (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-PRKCZ antibody	Rabbit polyclonal antibody to PKC zeta
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	75709226
NCBI GeneID	5590
NCBI Accession #	NP_001028753.1
NCBI GenBank Nucleotide #	NM_001033581.1
UniProt Accession #	Q05513; Q15207; Q5SYT5; Q969S4; A8K4N0; A8MU64; B7Z2J7; E9PCW2
Molecular Weight	56,136 Da
NCBI Official Full Name	protein kinase C zeta type isoform 2
NCBI Official Synonym Full Names	protein kinase C, zeta
NCBI Official Symbol	PRKCZ
NCBI Official Synonym Symbols	PKC2; PKC-ZETA
NCBI Protein Information	protein kinase C zeta type
UniProt Protein Name	Protein kinase C zeta type
UniProt Gene Name	PRKCZ
UniProt Synonym Gene Names	PKC2
UniProt Entry Name	KPCZ_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/polyclonal/pkc-zeta-antibody-immunoprecipitation-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/106138>