

Rabbit NOTCH1 Polyclonal Antibody | anti-NOTCH1 antibody

Anti-NOTCH1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA106146	Rabbit	Human, Mouse, Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

NOTCH1; Polyclonal Antibody; Anti-NOTCH1 Antibody; TAN1; Neurogenic locus notch homolog protein 1; Notch 1; hN1; Translocation-associated notch protein TAN-1; anti-NOTCH1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of NOTCH1 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	2,555
Applicable Applications for anti-NOTCH1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human NOTCH1. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of NOTCH1 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of NOTCH1 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of NOTCH1 expression in HeLa (A), NIH3T3 (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-NOTCH1 antibody

Rabbit polyclonal antibody to NOTCH1

NCBI AND UNIPROT INFORMATION

NCBI GI #	148833508
NCBI GeneID	4851
NCBI Accession #	NP_060087.3
NCBI GenBank Nucleotide #	NM_017617.3
UniProt Accession #	P46531; Q59ED8; Q5SXM3
Molecular Weight	272,505 Da
NCBI Official Full Name	neurogenic locus notch homolog protein 1 preproprotein
NCBI Official Synonym Full Names	notch 1
NCBI Official Symbol	NOTCH1
NCBI Official Synonym Symbols	hN1; AOS5; TAN1; AOVD1
NCBI Protein Information	neurogenic locus notch homolog protein 1
UniProt Protein Name	Neurogenic locus notch homolog protein 1
UniProt Gene Name	NOTCH1
UniProt Synonym Gene Names	TAN1; Notch 1; hN1; NEXT; NICD
UniProt Entry Name	NOTC1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/notch1-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/106146>