

Rabbit anti-Human MMP10 Polyclonal Antibody | anti-MMP10 antibody

Anti-MMP10 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA106148	Rabbit	Human	Flow Cytometry, Functional Assay, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MMP10; Polyclonal Antibody; Anti-MMP10 Antibody; STMY2; Stromelysin-2; SL-2; Matrix metalloproteinase-10; MMP-10; Transin-2; anti-MMP10 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of MMP10 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	476
Applicable Applications for anti-MMP10 antibody	FCM/FACS (Flow Cytometry), ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human MMP10. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of MMP10 staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of MMP10 staining in human prostate cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of MMP10 expression in Jurkat (A), HepG2 (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-MMP10 antibody

Rabbit polyclonal antibody to MMP10

NCBI AND UNIPROT INFORMATION

NCBI GI #	4505205
NCBI GeneID	4319
NCBI Accession #	NP_002416.1
NCBI GenBank Nucleotide #	NM_002425.2
UniProt Accession #	P09238; Q53HH9; B2R9X9
Molecular Weight	54,151 Da
NCBI Official Full Name	stromelysin-2 preproprotein
NCBI Official Synonym Full Names	matrix metallopeptidase 10
NCBI Official Symbol	MMP10
NCBI Official Synonym Symbols	SL-2; STMY2
NCBI Protein Information	stromelysin-2
UniProt Protein Name	Stromelysin-2
UniProt Gene Name	MMP10
UniProt Synonym Gene Names	STMY2; SL-2; MMP-10
UniProt Entry Name	MMP10_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/mmp10-antibody-flow-cytometry-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/106148>