

Rabbit MCL1 Polyclonal Antibody | anti-MCL1 antibody

MCL1 Polyclonal Antibody


CATALOG # AAA10630	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot
------------------------------	-----------------------	--	-------------------------------------

PRODUCT SYNONYMS


MCL1; Polyclonal Antibody; MCL1 Polyclonal Antibody; MCL1: MCL1; BCL2L3; EAT; MCL1L; MCL1S; MGC104264; MGC1839; Mcl-1; TM; anti-MCL1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	350
Applicable Applications for anti-MCL1 antibody	WB (Western Blot)
Application Notes	WB: 1:500 - 1:2000
Species	Human
Route	Synthetic Peptide
Immunogen	C term -peptide of human MCL1
Calculated Molecular Weight	37kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.




IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded mouse brain using MCL1 antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded mouse liver using MCL1 antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human gastric cancer using MCL1 antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human stomach using MCL1 antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human prostate using MCL1 antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human lung cancer using MCL1 antibody at dilution of 1:100 (40x lens).

ADDITIONAL INFORMATION

Related Product Information for anti-MCL1 antibody	Mcl-1 is an anti-apoptotic member of the Bcl-2 family originally isolated from the ML-1 human myeloid leukemia cell line during phorbol ester-induced differentiation along the monocyte/macrophage pathway (1). Similar to other Bcl-2 family members, Mcl-1 localizes to the mitochondria (2), interacts with and antagonizes pro-apoptotic Bcl-2 family members (3), and inhibits apoptosis induced by a number of cytotoxic stimuli (4). Mcl-1 differs from its other family members in its regulation at both the transcriptional and post-translational level. First, Mcl-1 has an extended amino-terminal PEST region, which is responsible for its relatively short half-life (1,2). Second, unlike other family members, Mcl-1 is rapidly transcribed via a PI3K/Akt dependent pathway, resulting in its increased expression during myeloid differentiation and cytokine stimulation (1,5-7). Mcl-1 is phosphorylated in response to treatment with phorbol ester, microtubule-damaging agents, oxidative stress, and cytokine withdrawal (8-11). Phosphorylation at Thr163, the conserved MAP kinase/ERK site located within the PEST region, slows Mcl-1 protein turnover (10) but may prime the GSK-3 mediated phosphorylation at Ser159 that leads to Mcl-1 destabilization (11). Mcl-1 deficiency in mice results in peri-implantation lethality (12). In addition, conditional disruption of the corresponding mcl-1 gene shows that Mcl-1 plays an important role in early lymphoid development and in the maintenance of mature lymphocytes (13).
Product Categories/Family for anti-MCL1 antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	83304396
NCBI GeneID	4170
UniProt Accession #	Q07820; Q9HD91; Q9NRQ3; Q9NRQ4; Q9UHR7; Q9UHR8; Q9UHR9; Q9UNJ1; B2R6B2; D3DV03; D3DV04
Molecular Weight	350
NCBI Official Full Name	Induced myeloid leukemia cell differentiation protein Mcl-1
NCBI Official Synonym Full Names	myeloid cell leukemia 1
NCBI Official Symbol	MCL1
NCBI Official Synonym Symbols	TM; EAT; MCL1L; MCL1S; Mcl-1; BCL2L3; MCL1-ES; bcl2-L-3; mcl1/EAT
NCBI Protein Information	induced myeloid leukemia cell differentiation protein Mcl-1; bcl-2-like protein 3; myeloid cell leukemia ES; bcl-2-related protein EAT/mcl1; myeloid cell leukemia sequence 1 (BCL2-related)
UniProt Protein Name	Induced myeloid leukemia cell differentiation protein Mcl-1
UniProt Gene Name	MCL1
UniProt Synonym Gene Names	BCL2L3; Bcl2-L-3
UniProt Entry Name	MCL1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/mcl1-antibody-western-blot-elisa/10630>