

## IgG (H+L) Secondary Antibody

HRP Rabbit Anti-Camel IgG (H+L) Antibody

|                              |                       |                                     |
|------------------------------|-----------------------|-------------------------------------|
| CATALOG #<br><b>AAA10762</b> | HOST<br><b>Rabbit</b> | APPLICATIONS<br><b>Western Blot</b> |
|------------------------------|-----------------------|-------------------------------------|


### PRODUCT SYNONYMS

IgG (H+L); HRP Rabbit Anti-Camel IgG (H+L) Antibody; HRP Rabbit Anti-Camel IgG (H+L): N/A; IgG (H+L) secondary antibody

### PRODUCT SPECIFICATIONS

|                                                          |                                                 |
|----------------------------------------------------------|-------------------------------------------------|
| Host                                                     | Rabbit                                          |
| Isotype                                                  | Horsedaish peroxidase conjugated IgG            |
| Purity/Purification                                      | Affinity Purification                           |
| Form/Format                                              | PBS with 0.01% thimerosal, 50% glycerol, pH7.3. |
| Applicable Applications for IgG (H+L) secondary antibody | WB (Western Blot)                               |
| Application Notes                                        | WB 1:500 - 1:1000                               |
| Immunogen                                                | Camel IgG                                       |
| Species Reactivity/ Cross-Reactivity                     | Llama                                           |
| Preparation and Storage                                  | Store at -20°C. Avoid freeze / thaw cycles.     |

### IMAGES



#### WB (Western Blot)

Western blot analysis of extracts of various cell lines, using HRP Rabbit Anti-Camelid VHH Antibody (AAA10762) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 3s.

ADDITIONAL INFORMATION

|                                                              |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|--------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Related Product Information for IgG (H+L) secondary antibody | Background: Secondary antibodies are affinity-purified antibodies which will work with target-specific primary antibody in the detection, sorting or purification of its specified target. Secondary antibodies offer increased versatility enabling users to use many detection systems (e.g. HRP, AP, fluorescence). They can also provide greater sensitivity through signal amplification as multiple secondary antibodies . Most commonly, secondary antibodies are generated by immunizing the host animal (different from host species of primary antibody) with a pooled population of normal immunoglobulins from the host species of primary antibody and can be further purified and modified (i.e. antibody fragmentation, label conjugation, etc.) to ensure well-characterized specificity to corresponding normal immunoglobulins. |
| Product Categories/Family for IgG (H+L) secondary antibody   | <a href="#">Secondary Antibody</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |

NCBI AND UNIPROT INFORMATION

|                  |                    |
|------------------|--------------------|
| Molecular Weight | Observed MW: 30kDa |
|------------------|--------------------|

**For Research Use Only.** Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | [contact@aaabiotech.com](mailto:contact@aaabiotech.com)  
Online datasheet: <https://stage.aaabiotech.com/rabbit-secondary-antibody/igg-hl/10762>