

Rabbit Proteasome 20S alpha 6 Polyclonal Antibody

Proteasome 20S alpha 6 antibody

CATALOG # AAA107907	HOST Rabbit	APPLICATIONS Western Blot, Immunohistochemistry, Immunocytochemistry, Immunofluorescence
-------------------------------	-----------------------	--


PRODUCT SYNONYMS

Proteasome 20S alpha 6; Polyclonal Antibody; Proteasome 20S alpha 6 antibody; Polyclonal Proteasome 20S alpha 6; Anti-Proteasome 20S alpha 6; 27 kDa prosomal protein; p27K; PROS27; multicatalytic endopeptidase complex iota chain; PSMA6; MGC22756; Proteasome S alpha 6 20; MGC2333; Proteasome S alpha 6-20; prosomal P27K protein; MGC23846; proteasome subunit alpha type-6; IOTA; macropain iota chain; anti-Proteasome 20S alpha 6 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Proteasome 20S alpha 6 antibody was purified by antigen-affinity chromatography
Form/Format	Supplied as a concentrated solution containing 0.1M Tris, 0.1M Glycine, 20% Glycerol (pH 7.0). 0.01% Thimerosal was added as a preservative.
Concentration	1 mg/ml (varies by lot)
Applicable Applications for anti-Proteasome 20S alpha 6 antibody	WB (Western Blot), IHC (Immunohistochemistry), ICC (Immunocytochemistry), IF (Immunofluorescence)
Biological Significance	The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit.
Cross-Reactivity	Human, Mouse
Preparation and Storage	Store at 4 degree C for short term storage. Aliquot and store at -20 degree C for long term storage. Avoid repeated freeze/thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemical staining of paraffin-embedded H1299 xenograft using Proteasome 20S alpha 6 antibody at a dilution of 1:100




IF (Immunofluorescence)

Immunofluorescence analysis of paraformaldehyde-fixed A431, using 20S Proteasome alpha6 antibody at 1:200 dilution.



WB (Western Blot)

Western blot analysis of 30 ug of whole cell lysate (A: 293T; B: A431; C: H1299; D: HeLa) using a 12 % SDS PAGE gel and Proteasome 20S alpha 6 antibody at a dilution of 1:1000



WB (Western Blot)

Western blot analysis of 50 ug of whole cell lysate (A: mouse brain) using a 12 % SDS PAGE gel and Proteasome 20S alpha 6 antibody at a dilution of 1:1000

ADDITIONAL INFORMATION

Related Product Information for anti-Proteasome 20S alpha 6 antibody	Affinity purified Rabbit polyclonal Proteasome 20S alpha 6 antibody
Product Categories/Family for anti-Proteasome 20S alpha 6 antibody	Proteases ; Inhibitors ; & Enzymes ; Purified Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

Molecular Weight	27 kDa
------------------	--------

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/proteasome-20s-alpha-6-antibody-western-blot-immunohistochemistry-immunocytochemistry-immunofluorescence-elisa/107907>