

Mouse HK1 Antibody | anti-HK1 antibody

HK1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA108770	Mouse	Human, Rat, Mouse	ELISA, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

HK1; HK1 Antibody; HKI; HXK1; HK1-ta; HK1-tb; HK1-tc; anti-HK1 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human, Rat, Mouse
Isotype	IgG1
Clone Number	7A7
Form/Format	Ascitic fluid containing 0.03% sodium azide.
Sequence Length	917
Applicable Applications for anti-HK1 antibody	ELISA, IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Purified recombinant fragment of human HK1 expressed in E. Coli.
Preparation and Storage	Store at 4 degree C, for long term storage, store at -20 degree C.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of NIH/3T3 cells using HK1 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human salivary gland tissues (left) and kidney tissues (right) using HK1 mouse mAb with DAB staining.



WB (Western Blot)

Western blot analysis using HK1 mouse mAb against Jurkat (1), HeLa (2), HepG2 (3), MCF-7 (4) and PC-12 (5) cell lysate.

ADDITIONAL INFORMATION

Related Product Information for anti-HK1 antibody	The hexokinases utilize Mg-ATP as a phosphoryl donor to catalyze the first step of intracellular glucose metabolism, the conversion of glucose to glucose-6-phosphate. Four hexokinase isoenzymes have been identified, including hexokinase I (HXK I), hexokinase II (HXK II), hexokinase III (HXK III) and hexokinase IV (HXK IV, also designated glucokinase or GCK). Hexokinases I-III each contain an N-terminal cluster of hydrophobic amino acids. Glucokinase lacks the N-terminal hydrophobic cluster. The hydrophobic cluster is thought to be necessary for membrane binding. This is substantiated by the finding that glucokinase has lower affinity for glucose than do the other hexokinases. HK I has been shown to be expressed in brain, kidney and heart tissues as well as in hepatoma cell lines.
Product Categories/Family for anti-HK1 antibody	Autophagy antibody ; Cancer ; Cardiovascular ; Cell Biology ; Epigenetics & Nuclear Signaling ; Developmental Biologys ; Immunology ; Drug Discovery Products ; Metabolism ; Neuroscience ; Signal Transduction ; Stem Cells

NCBI AND UNIPROT INFORMATION

NCBI GI #	188497754
NCBI GeneID	3098
NCBI Accession #	NP_000179.2
NCBI GenBank Nucleotide #	NM_000188.2
UniProt Accession #	P19367; O43443; O43444; O75574; Q5VTC3; Q96HC8; Q9NNZ4; Q9NNZ5; E9PCK0
Molecular Weight	102kDa
NCBI Official Full Name	hexokinase-1 isoform HKI
NCBI Official Synonym Full Names	hexokinase 1
NCBI Official Symbol	HK1
NCBI Official Synonym Symbols	HK; HKD; HKI; HXK1; RP79; HMSNR; HK1-ta; HK1-tb; HK1-tc; hexokinase
NCBI Protein Information	hexokinase-1
UniProt Protein Name	Hexokinase-1
UniProt Gene Name	HK1
UniProt Synonym Gene Names	HK I

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.