

Mouse anti-Human PDL2 Monoclonal Antibody | anti-PDL2 antibody

PDL2 Antibody [4E10]

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA10985	Mouse	Human	ELISA, Western Blot, Immunohistochemistry, Immunocytochemistry, Immunofluorescence, Flow Cytometry, Functional Assay

PRODUCT SYNONYMS

PDL2; Monoclonal Antibody; PDL2 Antibody [4E10]; B7DC; Btdc; CD273; PD-L2; PDCD1L2; bA574F11.2; Programmed cell death 1 ligand 2; Butyrophilin B7-DC; PD-1 ligand 2; programmed cell death 1 ligand; anti-PDL2 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Clone Number	4E10
Purity/Purification	PD-L2 Antibody is supplied as protein A purified IgG1.
Form/Format	Liquid; Pd-L2 Antibody is supplied in PBS containing 0.02% sodium azide and 50% glycerol.
Concentration	1 mg/mL (varies by lot)
Sequence Length	273
Applicable Applications for anti-PDL2 antibody	ELISA, WB (Western Blot), IHC (Immunohistochemistry), ICC (Immunocytochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Application Notes	<p>WB at 0.5-1 ug/mL IHC starting at 2 ug/mL IF start at 20 ug/mL.</p> <p>Antibody validated: Western Blot in human samples; Immunohistochemistry in human samples; Immunocytochemistry in human samples; Immunofluorescence in human samples and Flow Cytometry in mouse samples. All other applications and species not yet tested.</p>
Immunogen	PD-L2 antibody was raised against the extracellular domain of human PD-L2.
Conjugate	Unconjugated
Preparation and Storage	<p>Store at 4 degree C for three months and -20 degree C, stable for up to one year.</p> <p>As with all antibodies care should be taken to avoid repeated freeze thaw cycles.</p> <p>Antibodies should not be exposed to prolonged high temperatures.</p>

IMAGES



IF (Immunofluorescence)
Immunofluorescence of PD-L2 in human colon carcinoma tissue with PD-L2 antibody at 20 µg/mL. Green: PDL2 Antibody [4E10] (AAA10985) Blue: DAPI staining



FCM (Flow Cytometry)
Flow cytometry analysis of PD-L2 overexpressing HEK293 cells using PD-L2 antibody and control mouse IgG antibody at 10 µg/mL. Blue: Untransfected HEK293 cells. Yellow: PD-L2 overexpressing HEK293 cells.



IHC (Immunohistochemistry)
Immunohistochemistry of PD-L2 in human colon carcinoma tissue with PD-L2 antibody at 2 µg/mL.



IHC (Immunohistochemistry)
Immunohistochemistry of PD-L2 in human tonsil tissue with PD-L2 antibody at 2 µg/mL.



IF (Immunofluorescence)
Immunofluorescence of PD-L2 in human colon carcinoma tissue with PD-L2 antibody at 20 µg/mL. Green: PDL2 Antibody [4E10] Blue: DAPI staining



IF (Immunofluorescence)
Immunofluorescence of PD-L2 in human tonsil tissue with PD-L2 antibody at 20 µg/mL. Green: PDL2 Antibody [4E10] Blue: DAPI staining

ADDITIONAL INFORMATION

Related Product Information for anti-PDL2 antibody	Cell-mediated immune responses are initiated by T lymphocytes that are themselves stimulated by cognate peptides bound to MHC molecules on antigen-presenting cells (APC). T-cell activation is generally self-limited as activated T cells express receptors such as PD-1 (also known as PDCD-1) that mediate inhibitory signals from the APC. PD-1 can bind two different but related ligands, PD-L1 and PD-L2, both of which are thought to act as a negative regulator of T cell activation. However, it has been suggested that PD-L2 can act to stimulate an immunogenic response through an alternative receptor from PD-1.
Product Categories/Family for anti-PDL2 antibody	Immunology
References	Holling TM, Schooten E, and van Den Elsing PJ. Function and regulation of MHC class II molecules in T-lymphocytes: of mice and men. Hum. Immunol. 2004; 65:282-90. Ishida Y, Agata Y, Shibahara K, et al. Induced expression of PD-1, a novel member of the immunoglobulin gene superfamily, upon programmed cell death. EMBO J. 1992; 11:3887-95. LaGier J and Pober JS. Immune accessory functions of human endothelial cells are modulated by overexpression of B7-H1 (PDL1). Hum. Immunol. 2006; 67:568-78. Zhang Y, Chung Y, Bishop C, et al. Regulation of T cell activation and tolerance by PDL2. Proc. Natl. Acad. Sci. USA 2006; 103:11695-700.

NCBI AND UNIPROT INFORMATION

NCBI GI #	190014605
NCBI GeneID	80380
NCBI Accession #	NP_079515
NCBI GenBank Nucleotide #	NM_025239.3
UniProt Accession #	P9BQ51; Q14CN8; Q5T7Z6; Q6JXL8; Q6JXL9; Q9BQ51
Molecular Weight	Predicted: 30 kDa Observed: 38 kDa
NCBI Official Full Name	programmed cell death 1 ligand 2
NCBI Official Synonym Full Names	programmed cell death 1 ligand 2

NCBI Official Symbol	PDCD1LG2
NCBI Official Synonym Symbols	B7DC; Btdc; PDL2; CD273; PD-L2; PDCD1L2; bA574F11.2
NCBI Protein Information	programmed cell death 1 ligand 2
UniProt Protein Name	Programmed cell death 1 ligand 2
UniProt Gene Name	PDCD1LG2
UniProt Synonym Gene Names	B7DC; CD273; PDCD1L2; PDL2; PD-1 ligand 2; PD-L2; PDCD1 ligand 2; Programmed death ligand 2; B7-DC

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/pdl2-antibody-clone-4e10-western-blot-immunohistochemistry-immunocytochemistry-immunofluorescence-flow-cytometry-elisa/10985>