

Hexokinase-1 Recombinant Protein | HK1 recombinant protein

Recombinant Human Hexokinase-1

CATALOG # AAA113198	HOST E Coli or Yeast or Baculovirus or Mammalian Cell
-------------------------------	---

PRODUCT SYNONYMS

Hexokinase-1; Recombinant Human Hexokinase-1; Brain form hexokinaseHexokinase type I; HK I; HK1 recombinant protein

PRODUCT SPECIFICATIONS

Host	E Coli or Yeast or Baculovirus or Mammalian Cell
Purity/Purification	Greater or equal to 85% purity as determined by SDS-PAGE.
Form/Format	Lyophilized or liquid (Format to be determined during the manufacturing process)
Sequence Positions	1-917aa; Full Length Protein
Sequence	MIAAQLLAY FTELKDDQVK KIDKYLYAMR LSEDTLIDIM TRFRKEMKNG LSRDFNPTAT VKMLPTFVRS IPDGSEK GDF IALDLGGSSF RILRVQVNHE KNQNVHMESE VYDTPENIVH GSGSQLFDHV AECLGDFMEK RKIKDKKLPV GFTFSFPCQQ SKIDEAILIT WTKRFKASGV EGADVVKLLN KAIKKRGDYD ANIVAVVNDT VGTMMTCGYD DQHCEVGLII GTGTNACYME ELRHIDLVEG DEGRMCINTE WGAFGDDGSL EDIRTEFDRE IDRGS LNPGK QLFKEMVSGM YLGELVRLIL VKMAKEGLLF EGRITPELLT RGKFNTSDVS AIEKNKEGLH NAKEILTRLG VEPSDDDCVS VQHVCTIVSF RSANLVAATL GAILNRLRDN KGTPRLRTTV GVDGSLYKTH PQYSRRFHKT LRRLVPDSDV RFLLESSESG KGAAMVTAVA YRLAEQHRQI EETLAHFHLT KDMLEVKKR MRAEMELGLR KQTHNNNAV K MLPSFVRRTP DGTENGDFLA LDLGGTNFRV LLVKIRSGKK RTVEMHNKIY AIP EIMQGT GEELFDHIVS CISDFLDYMG IKGPRMPLGF TFSFPCQQT S LDAGILITWT KGFKATDCVG HDVVTLLRDA IKRREEFDLD VVAVVNDTVG TMMTCAYEEP TCEVGLIVGT GSNACYMEEM KNVEMVEGDQ GQMCINMEWG AFGDNGCLDD IRTHYDRLVD EYSLNAGKQR YEKMISGMYL GEIVRNILID FTKKGFLFRG QISETLKTRG IFETKFLSQI ESDRLALLQV RAILQQLGLN STCDD SILVK TVCGVVSRR A AQLCGAGMAA VVDKIRENRG LDRLNVTGVG DGTLYKLHPH FSRIMHQT VK ELSPKCNVSF LLS EDGSGKG AALITAVGVR LRTEASS
Preparation and Storage	Store at -20 degree C, for extended storage, conserve at -20 degree C or -80 degree C.

IMAGES



SDS-PAGE

ADDITIONAL INFORMATION

Product Categories/Family for HK1 recombinant protein	Metabolism
References	Human hexokinase sequences of amino- and carboxyl-terminal halves are homologous.Nishi S., Seino S., Bell G.I.Biochem. Biophys. Res. Commun. 157:937-943(1988) Structure of the human hexokinase type I gene and nucleotide sequence of the 5' flanking region.Ruzzo A., Andreoni F., Magnani M.Biochem. J. 331:607-613(1998) The DNA sequence and comparative analysis of human chromosome 10.Deloukas P., Earthrowl M.E., Grafham D.V., Rubenfield M., French L., Steward C.A., Sims S.K., Jones M.C., Searle S., Scott C., Howe K., Hunt S.E., Andrews T.D., Gilbert J.G.R., Swarbreck D., Ashurst J.L., Taylor A., Battles J., Bird C.P., Ainscough R., Almeida J.P., Ashwell R.I.S., Ambrose K.D., Babbage A.K., Bagguley C.L., Bailey J., Banerjee R., Bates K., Beasley H., Bray-Allen S., Brown A.J., Brown J.Y., Burford D.C., Burrill W., Burton J., Cahill P., Camire D., Carter N.P., Chapman J.C., Clark S.Y., Clarke G., Clee C.M., Clegg S., Corby N., Coulson A., Dhami P., Dutta I., Dunn M., Faulkner L., Frankish A., Frankland J.A., Garner P., Garnett J., Gribble S., Griffiths C., Grocock R., Gustafson E., Hammond S., Harley J.L., Hart E., Heath P.D., Ho T.P., Hopkins B., Horne J., Howden P.J., Huckle E., Hynds C., Johnson C., Johnson D., Kana A., Kay M., Kimberley A.M., Kershaw J.K., Kokkinaki M., Laird G.K., Lawlor S., Lee H.M., Leongamornlert D.A., Laird G., Lloyd C., Lloyd D.M., Loveland J., Lovell J., McLaren S., McLay K.E., McMurray A., Mashreghi-Mohammadi M., Matthews L., Milne S., Nickerson T., Nguyen M., Overton-Larty E., Palmer S.A., Pearce A.V., Peck A.I., Pelan S., Phillimore B., Porter K., Rice C.M., Rogosin A., Ross M.T., Sarafidou T., Sehra H.K., Shownkeen R., Skuce C.D., Smith M., Standring L., Sycamore N., Tester J., Thorpe A., Torcasso W., Tracey A., Tromans A., Tsolas J., Wall M., Walsh J., Wang H., Weinstock K., West A.P., Willey D.L., Whitehead S.L., Wilming L., Wray P.W., Young L., Chen Y., Lovering R.C., Moschonas N.K., Siebert R., Fechtel K., Bentley D., Durbin R.M., Hubbard T., Doucette-Stamm L., Beck S., Smith D.R., Rogers J.Nature 429:375-381(2004) Structure of the 5' region of the human hexokinase type I (HKI) gene and identification of an additional testis-specific HKI mRNA.Andreoni F., Ruzzo A., Magnani M.Biochim. Biophys. Acta 1493:19-26(2000) The erythrocyte-specific hexokinase isozyme (HKR) and the common hexokinase isozyme (HKI) are produced from a single gene by alternate promoters.Murakami K., Piomelli S.Blood 90:272-272(1998) Bienvenut W.V., Waridel P., Quadroni M.Submitted (MAR-2009) to UniProtKB Human hexokinase type I microheterogeneity is due to different amino-terminal sequences.Magnani M., Serafini G., Bianchi M., Casabianca A., Stocchi V.J. Biol. Chem. 266:502-505(1991) A recombinant human 'mini'-hexokinase is catalytically active and regulated by hexose 6-phosphates.Magnani M., Bianchi M., Casabianca A., Stocchi V., Daniele A., Altruda F., Ferrone M., Silengo L.Biochem. J. 285:193-199(1992) Identification of the cDNA for human red blood cell-specific hexokinase isozyme.Murakami K., Piomelli S.Blood 89:762-766(1997) Crystallization and preliminary X-ray analysis of human brain hexokinase.Aleshin A.E., Zeng C., Fromm H.J., Honatko R.B.FEBS Lett. 391:9-10(1996) Lys-N and trypsin cover complementary parts of the phosphoproteome in a refined SCX-based approach.Gauci S., Helbig A.O., Slijper M., Krijgsveld J., Heck A.J., Mohammed S.Anal. Chem. 81:4493-4501(2009) A mutation in an alternative untranslated exon of hexokinase 1 associated with hereditary motor and sensory neuropathy -- Russe (HMSNR) .Hantke J., Chandler D., King R., Wanders R.J., Angelicheva D., Tournev I., McNamara E., Kwa M., Guergueltcheva V., Kaneva R., Baas F., Kalaydjieva L.Eur. J. Hum. Genet. 17:1606-1614(2009) Initial characterization of the human central proteome.Burkard T.R., Planyavsky M., Kaupe I., Breitwieser F.P., Buerckstuemmer T., Bennett K.L., Superti-Furga G., Colinge J.BMC Syst. Biol. 5:17-17(2011) PKCepsilon promotes oncogenic functions of ATF2 in the nucleus while blocking its apoptotic function at mitochondria.Lau E., Kluger H., Varsano T., Lee K., Scheffler I., Rimm D.L., Ideker T., Ronai Z.A.Cell 148:543-555(2012) The mechanism of regulation of hexokinase new insights from the crystal structure of recombinant human brain hexokinase complexed with glucose and glucose-6-phosphate.Aleshin A.E., Zeng C., Bourenkov G.P., Bartunik H.D., Fromm H.J., Honzatko R.B.Structure 6:39-50(1998) Regulation of hexokinase I crystal structure of recombinant human brain hexokinase complexed with glucose and phosphate.Aleshin A.E., Zeng C., Bartunik H.D., Fromm H.J., Honzatko R.B.J. Mol. Biol. 282:345-357(1998) Binding of non-catalytic ATP to human hexokinase I highlights the structural components for enzyme-membrane association control.Rosano C., Sabini E., Rizzi M., Deriu D., Murshudov G., Bianchi M., Serafini G., Magnani M., Bolognesi M.Structure 7:1427-1437(1999) Crystal structures of mutant monomeric hexokinase I reveal multiple ADP binding sites and conformational changes relevant to allosteric regulation.Aleshin A.E., Kirby C., Liu X., Bourenkov G.P., Bartunik H.D., Fromm H.J., Honzatko R.B.J. Mol. Biol. 296:1001-1015(2000) Hexokinase mutations that produce nonspherocytic hemolytic anemia.Bianchi M., Magnani M.Blood Cells Mol. Dis. 21:2-8(1995) HK Utrecht missense mutation in the active site of human hexokinase associated with hexokinase deficiency and severe nonspherocytic hemolytic anemia.van Wijk R., Rijksen G., Huizinga E.G., Nieuwenhuis H.K., van Solinge W.W.Blood 101:345-347(2003)

NCBI AND UNIPROT INFORMATION

NCBI GI #	188497754
NCBI GeneID	3098

NCBI Accession #	NP_000179.2
NCBI GenBank Nucleotide #	NM_000188.2
UniProt Accession #	P19367 ; O43443 ; O43444 ; O75574 ; Q5VTC3 ; Q96HC8 ; Q9NNZ4 ; Q9NNZ5 ; E9PCK0
Molecular Weight	106.5 kDa
NCBI Official Full Name	hexokinase-1 isoform HKI
NCBI Official Synonym Full Names	hexokinase 1
NCBI Official Symbol	HK1
NCBI Official Synonym Symbols	HKD; HKI; HXK1; HMSNR; HK1-ta; HK1-tb; HK1-tc
NCBI Protein Information	hexokinase-1
UniProt Protein Name	Hexokinase-1
UniProt Gene Name	HK1
UniProt Synonym Gene Names	HK I
UniProt Entry Name	HXK1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/hk1-recombinant-protein-e-coli-or-yeast-or-baculovirus-or-mammalian-cell/113198>