

Collagen alpha-1(IV) chain Recombinant Protein | COL4A1 recombinant protein

Recombinant Human Collagen alpha-1(IV) chain protein

CATALOG # AAA113877	HOST E Coli or Yeast or Baculovirus or Mammalian Cell
-------------------------------	---


PRODUCT SYNONYMS

Collagen alpha-1(IV) chain; Recombinant Human Collagen alpha-1(IV) chain protein; COL4A1 recombinant protein

PRODUCT SPECIFICATIONS

Host	E Coli or Yeast or Baculovirus or Mammalian Cell
Purity/Purification	Greater or equal to 85% purity as determined by SDS-PAGE.
Form/Format	Lyophilized or liquid (Format to be determined during the manufacturing process)
Sequence Positions	1-362aa; Full-Length
Sequence	GCAGSGCGKC DCHGVKGQKG ERGLPGLQGV IGFPGMQGP GPQGGPGQKG DTGEPGLPGT KGTRGPPGAS GYPGNPGLPG IPGQDGPPGP PGIPGCNGTK GERGPLGPPG LPGFAGNPGP PGLPGMKGDP GEILGHVP
Preparation and Storage	Store at -20 degrees C. For long-term storage, store at -20 degrees C or -80 degrees C. Store working aliquots at 4 degrees C for up to one week. Repeated freezing and thawing is not recommended.

IMAGES



SDS-PAGE

ADDITIONAL INFORMATION

Related Product Information for COL4A1 recombinant protein	Type IV collagen is the major structural component of glomerular basement membranes (GBM), forming a 'chicken-wire' meshwork together with laminins, proteoglycans and entactin/nidogen. Arresten, comprising the C-terminal NC1 domain, inhibits angiogenesis and tumor formation. The C-terminal half is found to possess the anti-angiogenic activity. Specifically inhibits endothelial cell proliferation, migration and tube formation. Inhibits expression of hypoxia-inducible factor 1alpha and ERK1/2 and p38 MAPK activation. Ligand for alpha1/beta1 integrin.
Product Categories/Family for COL4A1 recombinant protein	Cancer

References

Structural organization of the gene for the alpha 1 chain of human type IV collagen. Soininen R., Huotari M., Ganguly A., Prockop D.J., Tryggvason K.J. *Biol. Chem.* 264:13565-13571(1989) The DNA sequence and analysis of human chromosome 13. Dunham A., Matthews L.H., Burton J., Ashurst J.L., Howe K.L., Ashcroft K.J., Beare D.M., Burford D.C., Hunt S.E., Griffiths-Jones S., Jones M.C., Keenan S.J., Oliver K., Scott C.E., Ainscough R., Almeida J.P., Ambrose K.D., Andrews D.T., Ashwell R.I.S., Babbage A.K., Bagguley C.L., Bailey J., Bannerjee R., Barlow K.F., Bates K., Beasley H., Bird C.P., Bray-Allen S., Brown A.J., Brown J.Y., Burrill W., Carder C., Carter N.P., Chapman J.C., Clamp M.E., Clark S.Y., Clarke G., Clee C.M., Clegg S.C., Cobley V., Collins J.E., Corby N., Coville G.J., Deloukas P., Dhami P., Dunham I., Dunn M., Earthrowl M.E., Ellington A.G., Faulkner L., Frankish A.G., Frankland J., French L., Garner P., Garnett J., Gilbert J.G.R., Gilson C.J., Ghorri J., Grafham D.V., Gribble S.M., Griffiths C., Hall R.E., Hammond S., Harley J.L., Hart E.A., Heath P.D., Howden P.J., Huckle E.J., Hunt P.J., Hunt A.R., Johnson C., Johnson D., Kay M., Kimberley A.M., King A., Laird G.K., Langford C.J., Lawlor S., Leongamornlert D.A., Lloyd D.M., Lloyd C., Loveland J.E., Lovell J., Martin S., Mashreghi-Mohammadi M., McLaren S.J., McMurray A., Milne S., Moore M.J.F., Nickerson T., Palmer S.A., Pearce A.V., Peck A.I., Pelan S., Phillimore B., Porter K.M., Rice C.M., Searle S., Sehra H.K., Shownkeen R., Skuce C.D., Smith M., Steward C.A., Sycamore N., Tester J., Thomas D.W., Tracey A., Tromans A., Tubby B., Wall M., Wallis J.M., West A.P., Whitehead S.L., Willey D.L., Wilming L., Wray P.W., Wright M.W., Young L., Coulson A., Durbin R.M., Hubbard T., Sulston J.E., Beck S., Bentley D.R., Rogers J., Ross M.T. *Nature* 428:522-528(2004) Completion of the amino acid sequence of the alpha 1 chain of human basement membrane collagen (type IV) reveals 21 non-triplet interruptions located within the collagenous domain. Brazel D., Oberbaeumer I., Dieringer H., Babel W., Glanville R.W., Deutzmann R., Kuehn K. *Eur. J. Biochem.* 168:529-536(1987) The structural genes for alpha 1 and alpha 2 chains of human type IV collagen are divergently encoded on opposite DNA strands and have an overlapping promoter region. Soininen R., Huotari M., Hostikka S.L., Prockop D.J., Tryggvason K.J. *Biol. Chem.* 263:17217-17220(1988) Amino acid sequence of the N-terminal aggregation and cross-linking region (7S domain) of the alpha 1 (IV) chain of human basement membrane collagen. Glanville R.W., Qian R.Q., Siebold B., Risteli J., Kuehn K. *Eur. J. Biochem.* 152:213-219(1985) Complete primary structure of the alpha 1-chain of human basement membrane (type IV) collagen. Soininen R., Haka-Risku T., Prockop D.J., Tryggvason K. *FEBS Lett.* 225:188-194(1987) Structure of human-basement-membrane (type IV) collagen. Complete amino-acid sequence of a 914-residue-long pepsin fragment from the alpha 1(IV) chain. Babel W., Glanville R.W. *Eur. J. Biochem.* 143:545-556(1984) cDNA clones coding for the pro-alpha1(IV) chain of human type IV procollagen reveal an unusual homology of amino acid sequences in two halves of the carboxyl-terminal domain. Pihlajaniemi T., Tryggvason K., Myers J.C., Kurkinen M., Lebo R., Cheung M.-C., Prockop D.J., Boyd C.D. *J. Biol. Chem.* 260:7681-7687(1985) Restricted homology between human alpha 1 type IV and other procollagen chains. Brinker J.M., Gudas L.J., Loidl H.R., Wang S.-Y., Rosenbloom J., Kefalides N.A., Myers J.C. *Proc. Natl. Acad. Sci. U.S.A.* 82:3649-3653(1985) The arrangement of intra- and intermolecular disulfide bonds in the carboxyterminal, non-collagenous aggregation and cross-linking domain of basement-membrane type IV collagen. Siebold B., Deutzmann R., Kuehn K. *Eur. J. Biochem.* 176:617-624(1988) Anti-angiogenic cues from vascular basement membrane collagen. Colorado P.C., Torre A., Kamphaus G., Maeshima Y., Hopfer H., Takahashi K., Volk R., Zamborsky E.D., Herman S., Sarkar P.K., Erickson M.B., Dhanabal M., Simons M., Post M., Kufe D.W., Weichselbaum R.R., Sukhatme V.P., Kalluri R. *Cancer Res.* 60:2520-2526(2000) Arresten, a collagen-derived inhibitor of angiogenesis. Fu J., Bai X., Wang W., Ruan C., Peng X., Yin B., Yuan J., Qiang B. *Molecular cloning and sequencing of human arresten gene.* Zheng Q.C., Song Z.F., Zheng Y.W., Li Y.Q., Shu X. *Zhonghua Shi Yan Wai Ke Za Zhi* 19:46-47(2002) Cloning and expression of arresten in *Escherichia coli* and *Pachia pastoris*. He A.B. *Construction of recombinant plasmid and prokaryotic expression in E. coli and biological activity analysis of human placenta arresten gene.* Zheng J.P., Tang H.Y., Chen X.J., Yu B.F., Xie J., Wu T.C. *Hepatobiliary Pancreat. Dis. Int.* 5:74-79(2006) Human alpha1 type IV collagen NC1 domain exhibits distinct antiangiogenic activity mediated by alpha1beta1 integrin. Sudhakar A., Nyberg P., Keshamouni V.G., Mannam A.P., Li J., Sugimoto H., Cosgrove D., Kalluri R. *J. Clin. Invest.* 115:2801-2810(2005) Characterization of the anti-angiogenic properties of arresten, an alpha1beta1 integrin-dependent collagen-derived tumor suppressor. Nyberg P., Xie L., Sugimoto H., Colorado P., Sund M., Holthaus K., Sudhakar A., Salo T., Kalluri R. *Exp. Cell Res.* 314:3292-3305(2008) An enzyme assisted RP-RPLC approach for in-depth analysis of human liver phosphoproteome. Bian Y., Song C., Cheng K., Dong M., Wang F., Huang J., Sun D., Wang L., Ye M., Zou H.J. *Proteomics* 96:253-262(2014) The 1.9-A crystal structure of the noncollagenous (NC1) domain of human placenta collagen IV shows stabilization via a novel type of covalent Met-Lys cross-link. Than M.E., Henrich S., Huber R., Ries A., Mann K., Kuhn K., Timpl R., Bourenkov G.P., Bartunik H.D., Bode W. *Proc. Natl. Acad. Sci. U.S.A.* 99:6607-6612(2002) Mutations in Col4a1 cause perinatal cerebral hemorrhage and porencephaly. Gould D.B., Phalan F.C., Breedveld G.J., van Mil S.E., Smith R.S., Schimenti J.C., Aguglia U., van der Knaap M.S., Heutink P., John S.W.M. *Science* 308:1167-1171(2005) Novel mutations in three families confirm a major role of COL4A1 in hereditary porencephaly. Breedveld G., de Coe I.F., Lequin M.H., Arts W.F.M., Heutink P., Gould D.B., John S.W.M., Oostra B., Mancini G.M.S. *J. Med. Genet.* 43:490-495(2006) Role of COL4A1 in small-vessel disease and hemorrhagic stroke. Gould D.B., Phalan F.C., van Mil S.E., Sundberg J.P., Vahedi K., Massin P., Bousser M.G., Heutink P., Miner J.H., Tournier-Lasserre E., John S.W.M. *N. Engl. J. Med.* 354:1489-1496(2006) COL4A1 mutation in Axenfeld-Rieger anomaly with leukoencephalopathy and stroke. Sibon I., Couprie Y., Menegon P., Bouchet J.P., Gorry P., Burgelin I., Calvas P., Orignac I., Dousset V., Lacombe D., Orgogozo J.M., Arveiler B., Goizet C. *Ann. Neurol.* 62:177-184(2007) COL4A1 mutations and hereditary angiopathy, nephropathy, aneurysms, and muscle cramps. Plaisier E., Gribouval O., Alamowitch S., Mougnot B., Prost C., Verpont M.C., Marro B., Desmettre T., Cohen S.Y., Rouillet E., Dracon M., Fardeau M., Van Agtmael T., Kerjaschki D., Antignac C., Ronco P.N. *Engl. J. Med.* 357:2687-2695(2007) A sulfilimine bond identified in collagen IV. Vanacore R., Ham A.-J.L., Voehler M., Sanders C.R., Conrads T.P., Veenstra T.D., Sharpless K.B., Dawson P.F., Hudson B.G. *Science* 325:1230-1234(2009) COL4A1 mutation in a patient with sporadic, recurrent intracerebral hemorrhage. Vahedi K., Kubis N., Boukobza M., Arnoult M., Massin P., Tournier-Lasserre E., Bousser M.G. *Stroke* 38:1461-1464(2007) COL4A1 mutation in two preterm siblings with antenatal onset of parenchymal

NCBI AND UNIPROT INFORMATION

NCBI GI #	734520332
NCBI GeneID	1282
NCBI Accession #	NP_001290039.1
NCBI GenBank Nucleotide #	NM_001303110.1
UniProt Accession #	P02462; Q1P9S9; Q5VWF6; Q86X41; Q8NF88; Q9NYC5; A7E2W4; B1AM70
Molecular Weight	39.9 kDa
NCBI Official Full Name	collagen alpha-1(IV) chain isoform 2
NCBI Official Synonym Full Names	collagen type IV alpha 1
NCBI Official Symbol	COL4A1
NCBI Official Synonym Symbols	BSVD; RATOR
NCBI Protein Information	collagen alpha-1(IV) chain
UniProt Protein Name	Collagen alpha-1(IV) chain
UniProt Gene Name	COL4A1
UniProt Entry Name	CO4A1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/col4a1-recombinant-protein-e-coli-or-yeast-or-baculovirus-or-mammalian-cell/113877>