

Human MIP-3 alpha/CCL20 ELISA Kit | CCL20 elisa kit

Human MIP-3 alpha/CCL20 PicoKine ELISA Kit

CATALOG #
AAA11575

REACTIVITY
Human


PRODUCT SYNONYMS

MIP-3 alpha/CCL20; Human MIP-3 alpha/CCL20 PicoKine ELISA Kit; C-C motif chemokine 20; chemokine (C-C motif) ligand 20; Beta chemokine exodus 1; Beta-chemokine exodus-1; C C motif chemokine 20; CC chemokine LARC; Ccl20; CCL20(2-70); CCL20_HUMAN; Chemokine (C C motif) ligand 20; CKb4; Exodus 1; LARC; Liver and activation regulated chemokine; Liver and activation-regulated chemokine; Macrophage inflammatory protein 3 alpha; MIP 3 alpha; MIP 3a; MIP-3-alpha; MIP3 alpha; MIP3A; MIP3alpha; SCYA20; Small inducible cytokine A20; Small inducible cytokine subfamily A (Cys Cys) member 20; Small-inducible cytokine A20; ST38; CCL20 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	Natural and recombinant Human CCL20
Sequence Length	96
Samples	Cell culture supernatants, cell lysates, serum and plasma (heparin, EDTA)
Detection Range	7.8 pg/ml - 500 pg/ml
Sensitivity	<1 pg/ml
Intra-assay Precision	Intra-Assay Precision (Precision within an assay): Three samples of known concentration were tested on one plate to assess intra-assay precision.
Inter-assay Precision	Inter-Assay Precision (Precision across assays): Three samples of known concentration were tested in separate assays to assess inter-assay precision.
Preparation and Storage	Store at 4 degree C for 6 months, at -20 degree C for 12 months. Avoid multiple freeze-thaw cycles (Shipped with wet ice.)

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CCL20 elisa kit	Background: Macrophage Inflammatory Protein 3alpha (MIP3alpha), also called Chemokine, cc motif, ligand 20 (CCL20). The MIP-3alpha/CCL20 gene was cloned and sequenced, revealing a four exon, three intron structure, and was localized by FISH analysis to 2q35-q36. MIP3alpha is predominantly expressed in lymph nodes, appendix, PBL, fetal liver, fetal lung and several cell lines. MIP3alpha/CCL20 and its receptor CCR6 are markedly upregulated in psoriasis, and they may play a role in the recruitment of T cells to lesional psoriatic skin. And Alanine MIP-3alpha and Serine MIP-3alpha, the two forms of MIP3alpha, that differ by one amino acid at the predicted signal peptide cleavage site. Both of them were chemically synthesized and tested for biological activity. And both flu antigen plus IL-2-activated CD4 (+) and CD8 (+) T lymphoblasts and cord blood-derived dendritic cells responded to Ser and Ala MIP-3alpha. Principle of the Assay: The Picokine™ Human CCL20 Pre-Coated ELISA (Enzyme-Linked Immunosorbent Assay) kit is a solid-phase immunoassay specially designed to measure Human CCL20 with a 96-well strip plate that is pre-coated with antibody specific for CCL20. The detection antibody is a biotinylated antibody specific for CCL20. The capture antibody is monoclonal antibody from mouse and the detection antibody is polyclonal antibody from goat. The kit contains recombinant Human CCL20 with immunogen: Expression system for standard: E.coli; Immunogen sequence: A27-M96. The kit is analytically validated with ready-to-use reagents. To measure Human CCL20, add standards and samples to the wells, then add the biotinylated detection antibody. Wash the wells with PBS or TBS buffer, and add Avidin-Biotin-Peroxidase Complex (ABC-HRP). Wash away the unbounded ABC-HRP with PBS or TBS buffer and add TMB. TMB is an HRP substrate and will be catalyzed to produce a blue color product, which changes into yellow after adding the acidic stop solution. The absorbance of the yellow product at 450nm is linearly proportional to Human CCL20 in the sample. Read the absorbance of the yellow product in each well using a plate reader, and benchmark the sample wells' readings against the standard curve to determine the concentration of Human CCL20 in the sample.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	2829668
NCBI GeneID	6364
UniProt Accession #	P78556; Q53S51; Q99664
Molecular Weight	10,691 Da
NCBI Official Full Name	C-C motif chemokine 20
NCBI Official Synonym Full Names	chemokine (C-C motif) ligand 20
NCBI Official Symbol	CCL20
NCBI Official Synonym Symbols	CKb4; LARC; ST38; MIP3A; Exodus; MIP-3a; SCYA20; MIP-3-alpha
NCBI Protein Information	C-C motif chemokine 20; CC chemokine LARC; beta chemokine exodus-1; small-inducible cytokine A20; macrophage inflammatory protein 3 alpha; liver and activation-regulated chemokine; small inducible cytokine subfamily A (Cys-Cys), member 20
UniProt Protein Name	C-C motif chemokine 20
UniProt Gene Name	CCL20
UniProt Synonym Gene Names	LARC; MIP3A; SCYA20; MIP-3-alpha
UniProt Entry Name	CCL20_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/mip-3-alpha-ccl20/11575>