

Rabbit anti-Human Tumor necrosis factor receptor type 1-associated DEATH domain protein Polyclonal Antibody | anti-TRADD antibody

Rabbit anti-human Tumor necrosis factor receptor type 1-associated DEATH domain protein polyclonal Antibody(TRADD)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA119074	Rabbit	Human	Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

Tumor necrosis factor receptor type 1-associated DEATH domain protein; Polyclonal Antibody; Rabbit anti-human Tumor necrosis factor receptor type 1-associated DEATH domain protein polyclonal Antibody(TRADD); TNFR1-associated DEATH domain protein; TNFRSF1A-associated via death domain; TRADD; anti-TRADD antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purified
Sequence Length	312
Applicable Applications for anti-TRADD antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Conjugate	Non-conjugated
Immunogen	Recombinant human Tumor necrosis factor receptor type 1-associated DEATH domain protein (1-312aa)
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-1163 / sc-7868
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human colon cancer using AAA119074 at dilution 1:100



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human endometrial cancer using AAA119074 at dilution 1:100



WB (Western Blot)

All lanes: Tumor necrosis factor receptor type 1-associated DEATH domain protein antibody at 12ug/ml+ MCF-7 whole cell lysate. Secondary Goat polyclonal to rabbit at 1/10000 dilution. Predicted band size: 35,28 kDa. Observed band size: 34 kDa

ADDITIONAL INFORMATION

Related Product Information for anti-TRADD antibody	The nuclear form acts as a tumor suppressor by preventing ubiquitination and degradation of isoform p19ARF/ARF of CDKN2A by TRIP12: acts by interacting with TRIP12, leading to disrupt interaction between TRIP12 and isoform p19ARF/ARF of CDKN2A (By similarity). Adapter molecule for TNFRSF1A/TNFR1 that specifically associates with the cytoplasmic domain of activated TNFRSF1A/TNFR1 mediating its interaction with FADD. Overexpression of TRADD leads to two major TNF-induced responses, apoptosis and activation of NF-kappa-B.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	13378137
NCBI GeneID	8717
NCBI Accession #	NP_003780.1
NCBI GenBank Nucleotide #	NM_003789.3
UniProt Accession #	Q15628; Q52NZ1; B2RDS3; B3KQZ9
Molecular Weight	27,904 Da
NCBI Official Full Name	tumor necrosis factor receptor type 1-associated DEATH domain protein
NCBI Official Synonym Full Names	TNFRSF1A-associated via death domain
NCBI Official Symbol	TRADD
NCBI Official Synonym Symbols	Hs.89862
NCBI Protein Information	tumor necrosis factor receptor type 1-associated DEATH domain protein
UniProt Protein Name	Tumor necrosis factor receptor type 1-associated DEATH domain protein
UniProt Gene Name	TRADD
UniProt Synonym Gene Names	TNFR1-associated DEATH domain protein
UniProt Entry Name	TRADD_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/tumor-necrosis-factor-receptor-type-1-associated-death-domain-protein-antibody-immunohistochemistry-western-blot-elisa/119074>